herpy

PIJLGIFKIKKERS

Blue-bellied Poison Frog

Andinobates minutus

Andinobates minutus is one of the smallest poison dart frogs, only 13-16 mm from snout to vent. It lives in the leaf litter of humid lowland forest from central Panama to the Pacific coast of Colombia, and the male carries the tadpoles up into bromeliads. The species is not collected for the pet trade and is hardly kept, so little is known about its care.

Familie: Dendrobatidae Vroegere naam: *Ranitomeya minuta; Dendrobates minutus; Minyobates minutus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**13-16 mm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**1-8**
Larvenperiode	**60-80 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kant-en-klaar voer**
Broedzorg	**Draagt de kikkervisjes**

VERZORGING

Min. terrarium	**45 × 45 × 45 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**22-25 °C**
Temperatuur ’s nachts	**19-22 °C**
Luchtvochtigheid	**80-100%**
Water	**Geen waterbak**
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Andinobates minutus*. The care is based on the habitat: warm, humid lowland forest with rain for most of the year and a thick layer of leaf litter.
TEMPERATUUR	The daytime temperature should be 22-25 °C and the night 19-22 °C. The temperature must not exceed 27 °C for long, and above 29 °C is dangerous, so the enclosure must not stand in sun or close to a heat source.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 80-100 %. Spray the enclosure 1-2 times a day with rainwater or reverse osmosis water so that the leaf litter is kept damp, or use a rain system on a timer.
WATER EN WATERKWALITEIT	The adult frogs do not need a water bowl; they take up water through the skin from spraying and the damp leaf litter.
VERLICHTING EN UVB	The species is diurnal and lives in shade on the forest floor. LED lighting for about 12 hours a day is enough for the plants. UVB is not necessary; a weak UVB light in Ferguson zone 1 can be used if there is plenty of shade.
VOEDING	In the wild *Andinobates minutus* lives on mites, ants, springtails and insect larvae.
GEDRAG EN OMGANG	The species is diurnal and stays in the leaf litter and by the hiding places. The males call from fixed spots and defend their territory, so a pair is the way to keep them.
NATUURLIJK VERSPREIDINGSGEBIED	Colombia en Panama.