herpy

PIJLGIFKIKKERS

Andean Poison Frog

Andinobates opisthomelas

Andinobates opisthomelas is a small, diurnal poison dart frog of 15-20 mm from the Andes of Colombia, where it lives in the leaf litter of humid mountain forest in Antioquia and Caldas. The species is Vulnerable in the wild, and almost nothing is known about its care.

Familie: Dendrobatidae Vroegere naam: *Ranitomeya opisthomelas; Dendrobates opisthomelas; Minyobates opisthomelas*

KERNGEGEVENS

HET DIER

Volwassen grootte	**15-20 mm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Bergbos**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**1-2**
Uitkomen	**16-24 dagen**
Larvenperiode	**108-112 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Draagt de kikkervisjes**

VERZORGING

Min. terrarium	**45 × 50 × 60 cm** L × B × H voor 5 volwassen dieren
Temperatuur overdag	**22-25 °C**
Temperatuur ’s nachts	**18-21 °C**
Luchtvochtigheid	**80-100%**
Water	**Geen waterbak**
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Kwetsbaar (VU)**

Verzorging in het kort

HET TERRARIUM	Almost nothing is known about the care of *Andinobates opisthomelas*, and the species is hardly kept. The care is based on the habitat, the cool and humid mountain forest with leaf litter and bromeliads, and on close relatives.
TEMPERATUUR	The daytime temperature should be 22-25 °C, and at night it may fall to 18-21 °C as in the mountain forest. Keep the temperature below 27 °C.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 80-100 % and never below 70 %. Spray twice a day, morning and afternoon, or use a fogger or rain system with reverse osmosis water or rainwater.
WATER EN WATERKWALITEIT	A water bowl is not needed; the frogs drink droplets from plants and spray water. For the tadpoles, use bromeliads with water in the leaf axils or small plastic cups with 2-3 cm of water.
VERLICHTING EN UVB	The light is primarily for the plants: plant lighting for 10-12 hours a day provides a daily rhythm and growth.
VOEDING	The food is small arthropods: springtails, small fruit flies (*Drosophila melanogaster*) and small isopods. Newly metamorphosed froglets take only springtails.
GEDRAG EN OMGANG	*Andinobates opisthomelas* is diurnal and lives on the ground, so it stays low in the enclosure, in the leaf litter and by the bromeliads.
NATUURLIJK VERSPREIDINGSGEBIED	Colombia.