herpy

PIJLGIFKIKKERS

Anthony's Poison Arrow Frog

Epipedobates anthonyi

Epipedobates anthonyi is a small, red and striped poison frog of 19-27 mm from south-western Ecuador and north-western Peru. It is the frog that has been sold for decades as "tricolor", and it is probably the most widely kept and bred poison frog of all: hardy, bold, vocal and a prolific breeder.

Familie: Dendrobatidae Vroegere naam: *Dendrobates anthonyi; Phyllobates anthonyi*

KERNGEGEVENS

HET DIER

Volwassen grootte	**19-27 mm** kop-romplengte
Levensduur	**5-14 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Droog bos**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Geslachtsrijp	**5-10 maanden**
Eieren per legsel	**10-40**
Uitkomen	**9-14 dagen** bij 22-24 °C
Larvenperiode	**43-89 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**40 × 40 × 40 cm** L × B × H voor 3 volwassen dieren
Temperatuur overdag	**22-26 °C**
Temperatuur ‘s nachts	**18-21 °C**
Luchtvochtigheid	**80-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Gevoelig (NT)**

Verzorging in het kort

HET TERRARIUM	A terrarium of at least 40 × 40 × 40 cm suits one male and two females of *Epipedobates anthonyi*. A group of 5-10 animals needs at least 60 × 60 × 40 cm, and more animals need more floor area.
TEMPERATUUR	Keep 22-26 °C by day with both cooler and warmer zones, and let the temperature fall to room temperature at night, around 18-21 °C. The temperature must not exceed 27 °C for long, and above 29 °C is dangerous.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 80-100 %. The species tolerates down to 50 % for short periods if it has access to water, but low humidity without water can quickly become fatal.
WATER EN WATERKWALITEIT	Make one or more small water holes with about 2 cm of water, e.g. saucers, drinking dishes or half Brazil nut shells that the frogs can easily climb out of.
VERLICHTING EN UVB	The light is for the plants. Fluorescent tubes or LED bars for about 12 hours a day are enough, and the day length may vary a little over the year.
VOEDING	*Epipedobates anthonyi* is small but greedy and takes relatively large prey.
GEDRAG EN OMGANG	*Epipedobates anthonyi* is diurnal, bold and often out in the open, and the males call with a trill, mostly in the morning and after misting.
NATUURLIJK VERSPREIDINGSGEBIED	Ecuador en Peru.