herpy

PIJLGIFKIKKERS

Marbled Poison Frog

Epipedobates boulengeri

Epipedobates boulengeri is a small, diurnal poison dart frog of about 17-24 mm with a dark brown back and an unbroken, pale stripe along the side. It lives in the lowland rainforest along the Pacific coast from south-western Colombia to north-westernmost Ecuador. The species is hardly ever kept, and little is known about its care.

Familie: Dendrobatidae Vroegere naam: *Dendrobates boulengeri; Phyllobates boulengeri*

KERNGEGEVENS

HET DIER

Volwassen grootte	**17-24 mm** kop-romplengte
Levensduur	**8-10 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**10-30**
Uitkomen	**9-14 dagen**
Larvenperiode	**42-70 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**40 × 40 × 40 cm** L × B × H voor 3 volwassen dieren
Temperatuur overdag	**22-26 °C**
Temperatuur 's nachts	**19-22 °C**
Luchtvochtigheid	**80-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Epipedobates boulengeri*, and the species is hardly ever kept.
TEMPERATUUR	Keep 22-26 °C during the day with slightly cooler and slightly warmer spots in the enclosure, and 19-22 °C at night. This is normally achieved with room temperature and the lighting; heat mats are unnecessary.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 80-100 %. Spray several times a day or use a rain system, for example three times a day for 20-60 seconds, preferably supplemented with a fogger.
WATER EN WATERKWALITEIT	A shallow water bowl with 1-2 cm of water, for example a low saucer or half a Brazil nut shell, gives the frogs a place to drink and bathe and is a breeding site when the male is to deposit tadpoles.
VERLICHTING EN UVB	The light is for the plants: fluorescent tubes or LED bars for about 12 hours a day. The closely related species thrive without UVB, so it is optional.
VOEDING	The frogs are given live, small prey: springtails and fruit flies (*Drosophila melanogaster*), and adults also take the larger *Drosophila hydei* and newly hatched crickets.
GEDRAG EN OMGANG	The species of the genus are diurnal, ground-dwelling and territorial, and the males call from fixed spots. The call has not been described in more detail for *E. boulengeri* in the narrow sense.
NATUURLIJK VERSPREIDINGSGEBIED	Colombia en Ecuador.