herpy

PIJLGIFKIKKERS

Darwin Wallace poison frog

Epipedobates darwinwallacei

Epipedobates darwinwallacei is a small, black poison dart frog of 13-20 mm with orange to yellow spots. It lives in the montane forest around Mindo in north-western Ecuador and is threatened. Since 2024 it has been regarded as a synonym of *Epipedobates espinosai*. Little is known about its care.

Familie: Dendrobatidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**13-20 mm** kop-romplengte
Levensduur	**8-10 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud en bergbos**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**10-30**
Uitkomen	**9-14 dagen**
Larvenperiode	**42-70 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**40 × 40 × 40 cm** L × B × H voor 3 volwassen dieren
Temperatuur overdag	**20-24 °C**
Temperatuur ‘s nachts	**18-20 °C**
Luchtvochtigheid	**80-100%**
Water	**Watergedeelte** diepte 2-5 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Bedreigd (EN)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Epipedobates darwinwallacei*, and the species is hardly ever kept.
TEMPERATUUR	Keep 20-24 °C during the day and 18-20 °C at night. This corresponds to the cool end of what suits *Epipedobates tricolor* from the same elevation. It is normally achieved with room temperature and the lighting.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 80-100 %. Spray several times a day or use a rain system, for example three times a day for 20-60 seconds, preferably supplemented with a fogger.
WATER EN WATERKWALITEIT	Provide a small, permanent water feature with 2-5 cm of water and a gentle slope or ramp so that the frogs can easily climb out.
VERLICHTING EN UVB	The light is for the plants: fluorescent tubes or LED bars for about 12 hours a day. The closely related species thrive without UVB, so it is optional.
VOEDING	The frogs are given live, small prey: springtails and fruit flies (*Drosophila melanogaster*), and adults also take the larger *Drosophila hydei* and newly hatched crickets.
GEDRAG EN OMGANG	*E. darwinwallacei* is diurnal and moves openly between grass and leaf litter. The call is very similar to the call of the brown *E. espinosai* from the lowlands. The males of the genus are territorial and call from fixed spots.
NATUURLIJK VERSPREIDINGSGEBIED	Ecuador.