herpy

PIJLGIFKIKKERS

Rio Santiago Poison Frog

Excidobates captivus

Excidobates captivus is a very small, diurnal poison frog of 12-17 mm from the border region between Peru and Ecuador. It is black with orange-red spots in rows along the back and yellow spots on the limbs and belly, and it lives on the forest floor in humid rainforest and montane forest.

Familie: Dendrobatidae Vroegere naam: *Dendrobates captivus; Ranitomeya captiva; Adelphobates captivus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**12-17 mm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud en bergbos**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**4-8**
Larvenperiode	**65-75 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Draagt de kikkervisjes**

VERZORGING

Min. terrarium	**40 × 40 × 40 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**22-26 °C**
Temperatuur ’s nachts	**18-22 °C**
Luchtvochtigheid	**80-100%**
Water	**Geen waterbak**
UVB	**Optioneel**
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Kwetsbaar (VU)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Excidobates captivus*. The species lives on the forest floor in warm to mild, permanently humid forest.
TEMPERATUUR	The temperature at the species' localities has not been measured. 22-26 °C by day and 18-22 °C at night lies between the climate in the lowlands and at 800-1200 m elevation.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 80-100 %, and the substrate should be damp all year. Mist with reverse-osmosis or rainwater once or twice a day, or use a rain or fogging system. Make sure the air does not stand still.
WATER EN WATERKWALITEIT	A water dish is not necessary, because the frogs get moisture from the misting and the damp leaf litter. The water in bromeliads and film canisters is replaced with fresh reverse-osmosis or rainwater every week.
VERLICHTING EN UVB	The species lives in shade on the forest floor, so the lighting should first and foremost support plant growth and a day of 12 hours. UVB is not necessary, but optional.
VOEDING	*Excidobates captivus* is so small that it can only take very small feeder insects: springtails, flightless fruit flies (*Drosophila melanogaster*) and the smallest woodlice.
GEDRAG EN OMGANG	The species is diurnal, but hides a great deal in the foliage. The male calls short, shrill sounds from hidden spots. It is best kept as a pair, because several animals together can cause conflict.
NATUURLIJK VERSPREIDINGSGEBIED	Peru en Ecuador.