herpy

PIJLGIFKIKKERS

Trinidad Poison Frog

Mannophryne trinitatis

Mannophryne trinitatis is a small, brown, diurnal frog of 25-28 mm that occurs only on Trinidad, where it lives by clear, stony streams in humid forest. The male can turn jet black when calling, and the female has a clear yellow throat. The species is hardly ever kept, and little is known about its care.

Familie: Aromobatidae Vroegere naam: *Phyllobates trinitatis*

KERNGEGEVENS

HET DIER

Volwassen grootte	**25-28 mm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud en moerasgebied**
Huidgif	**Geen bekend**
Moeilijkheidsgraad	**Ervaren**

VERZORGING

Min. terrarium	**60 × 40 × 45 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**23-27 °C**
Temperatuur 's nachts	**20-22 °C**
Luchtvochtigheid	**80-100%**
Water	**Watergedeelte** diepte 3-5 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**6-13**
Uitkomen	**21 dagen**
Larvenperiode	**40-50 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little is known about keeping *Mannophryne trinitatis*.
TEMPERATUUR	The daytime temperature should be 23-27 °C and the night 20-22 °C. Frogs from forest streams tolerate heat poorly, so keep the temperature below 28 °C.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 80-100 %. Mist daily so that leaves and stones are wet. A fogging system makes it easier to keep the humidity stable. Provide ventilation so that the air does not become stagnant.
WATER EN WATERKWALITEIT	The species needs a permanent water section with shallow, clean and preferably gently flowing water, 3-5 cm deep, with stones and leaves.
VERLICHTING EN UVB	UVB is hardly necessary, but a low dose (Ferguson zone 1) is optional. Plant lighting for 10-12 hours keeps moss and ferns alive. Provide shade with plants and bark so that the frogs can move away from the bright light.
VOEDING	The species needs small arthropods: small and large fruit flies (*Drosophila melanogaster* and *Drosophila hydei*), springtails and small isopods.
GEDRAG EN OMGANG	*Mannophryne trinitatis* is diurnal and territorial. Females defend their spots more often than males, with threats, wrestling and chasing, and they display the yellow throat as a threat.
NATUURLIJK VERSPREIDINGSGEBIED	Trinidad en Tobago.