herpy

PIJLGIFKIKKERS

Polkadot Poison Frog

Oophaga arborea

Oophaga arborea is a small poison dart frog of 20-22 mm that lives high up in the trees in the humid montane forest of western Panama. It is black or brown with round yellow spots, breeds in bromeliads 10-16 m above the ground, and the tadpoles feed only on eggs from the female. The species is critically endangered.

Familie: Dendrobatidae Vroegere naam: *Dendrobates arboreus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**20-22 mm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend**
Leefgebied	**Bergbos**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Expert**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**In bladoksels (bijv. bromelia's)**
Eieren per legsel	**4-8**
Uitkomen	**14 dagen**
Larvenperiode	**40-50 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Onbevruchte eieren van het vrouwtje**
Broedzorg	**Bewaakt de eieren, draagt de kikkervisjes en voedt de kikkervisjes met eieren**

VERZORGING

Min. terrarium	**60 × 45 × 80 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**20-24 °C**
Temperatuur 's nachts	**15-19 °C**
Luchtvochtigheid	**70-100%**
Water	**Geen waterbak**
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Ernstig bedreigd (CR)**

Verzorging in het kort

HET TERRARIUM	Almost nothing is known about the care of *Oophaga arborea*, and the species is almost unknown in captivity.
TEMPERATUUR	The daytime temperature should be 20-24 °C, and at night it should fall by at least 4-5 °C, to about 15-19 °C. This corresponds to the climate of the cloud forest with an annual mean temperature of about 19 °C.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 70-80 % by day and rise to 100 % at night. Use plenty of mist and rain, and let the enclosure dry quickly afterwards with the help of a fan; stagnant, wet air is dangerous.
WATER EN WATERKWALITEIT	A water bowl is not necessary. The frogs get moisture from mist, rain and the water-filled bromeliads, which are also used for the tadpoles.
VERLICHTING EN UVB	The species lives high up in the canopy, where the light is stronger than on the forest floor, but the mist dampens it.
VOEDING	Adults are fed small, flightless fruit flies, springtails, newly hatched micro crickets and field plankton; the same feeder insects are used for the other species of the genus.
GEDRAG EN OMGANG	*Oophaga arborea* is a diurnal tree frog that is rarely seen on the ground.
NATUURLIJK VERSPREIDINGSGEBIED	Panama.