herpy

PIJLGIFKIKKERS

Granular Poison Frog

Oophaga granulifera

Oophaga granulifera is a small, diurnal poison dart frog of 19-22 mm from the Pacific side of Costa Rica and the far west of Panama. The skin is distinctly granular, the back red, orange or olive-green, and the legs and belly blue-green to turquoise. The species lives along shaded forest streams, and the female feeds her tadpoles with unfertilised eggs.

Familie: Dendrobatidae Vroegere naam: *Dendrobates granuliferus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**19-22 mm** kop-romplengte
Levensduur	**6 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

VERZORGING

Min. terrarium	**80 × 40 × 40 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**21-25 °C**
Temperatuur ’s nachts	**17-21 °C**
Luchtvochtigheid	**80-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Geslachtsrijp	**12-18 maanden**
Eieren per legsel	**2-5**
Uitkomen	**19-20 dagen**
Larvenperiode	**87-212 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Onbevruchte eieren van het vrouwtje**
Broedzorg	**Bewaakt de eieren, draagt de kikkervisjes en voedt de kikkervisjes met eieren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Kwetsbaar (VU)**

Verzorging in het kort

HET TERRARIUM	Little is known about the husbandry of *Oophaga granulifera*, and the species is rare in the hobby.
TEMPERATUUR	The daytime temperature should be 21-25 °C, and the frogs are most active at 20-22 °C. At night the temperature should fall by 4-5 °C, to about 17-21 °C. Seasonal variation is minimal, about 1-2 °C.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 80-90 %, and 100 % early in the morning and late in the day when the enclosure is misted.
WATER EN WATERKWALITEIT	A shallow water dish of 1-2 cm depth, for example a low bowl sunk into the substrate or a water section between the stones, provides drinking water and humidity.
VERLICHTING EN UVB	The species lives on the shady forest floor, so the light must first and foremost cover the plants' needs.
VOEDING	Adults are fed small, flightless fruit flies, springtails, newly hatched micro crickets, the smallest waxworm larvae and field plankton.
GEDRAG EN OMGANG	*Oophaga granulifera* is shy and lives very secretively, especially in the dry season, when it almost completely shuts down.
NATUURLIJK VERSPREIDINGSGEBIED	Costa Rica en Panama.