herpy

PIJLGIFKIKKERS

Harlequin Poison Frog

Oophaga histrionica

Oophaga histrionica is a diurnal poison frog of 3-4 cm from the rainforest of western Colombia, brown to black with large orange spots. The tadpoles live exclusively on infertile eggs that the female lays for them, which makes the species one of the most difficult poison frogs to breed. It is critically endangered, and the delimitation of the species was changed in 2018.

Familie: Dendrobatidae Vroegere naam: *Dendrobates histrionicus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**3-4 cm** kop-romplengte
Levensduur	**9 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Expert**

VERZORGING

Min. terrarium	**50 × 50 × 50 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**24-26 °C**
Temperatuur 's nachts	**18-21 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Geslachtsrijp	**12 maanden**
Eieren per legsel	**4-20**
Uitkomen	**10 dagen**
Larvenperiode	**90 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Onbevruchte eieren van het vrouwtje**
Broedzorg	**Bewaakt de eieren, draagt de kikkervisjes en voedt de kikkervisjes met eieren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Ernstig bedreigd (CR)**

Verzorging in het kort

HET TERRARIUM	A pair needs a rainforest vivarium of at least 50 × 50 × 50 cm, and larger is better. *Oophaga* species can be territorial, so keep only one pair unless the vivarium is very large.
TEMPERATUUR	The daytime temperature should be 24-26 °C, and at night it is lowered by 5-6 degrees, to 18-21 °C. The temperature only needs to vary by 1-2 degrees over the year, and there is no winter rest.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 70-80 % in the middle of the day and up to 100 % morning and evening. This is easiest to achieve with a rain system and a fogging system on timers.
WATER EN WATERKWALITEIT	A shallow water dish with about 1 cm of water and a ramp of stones or moss is enough for the adults. Use reverse-osmosis water, rain water or dechlorinated tap water, and change the water often.
VERLICHTING EN UVB	The lighting is first and foremost for the bromeliads and the other plants. Provide about 12 hours of light a day with LED or fluorescent tubes that do not give off much heat.
VOEDING	*Oophaga histrionica* eats small arthropods, in the wild especially ants, termites, mites and small beetles.
GEDRAG EN OMGANG	*Oophaga histrionica* is diurnal. The male calls from a low perch, and males defend their territories. Therefore keep only one pair in the enclosure. Do not handle the frogs.
NATUURLIJK VERSPREIDINGSGEBIED	Colombia.