herpy

PIJLGIFKIKKERS

Lehmann's Poison Frog

Oophaga lehmanni

Oophaga lehmanni is a rare, diurnal poison frog from a small area in western Colombia, black or dark brown with broad bands of red, orange or yellow. It is critically endangered in the wild, and only captive-bred animals belong in an enclosure. The species requires an experienced keeper: it does not tolerate heat, and the tadpoles live exclusively on unfertilised eggs from the female.

Familie: Dendrobatidae Vroegere naam: *Dendrobates lehmanni*

KERNGEGEVENS

HET DIER

Volwassen grootte	**3-4 cm** kop-romplengte
Levensduur	**4-15 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud en bergbos**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op bladeren boven het water**
Eieren per legsel	**3-10**
Uitkomen	**14-28 dagen**
Larvenperiode	**60-90 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Onbevruchte eieren van het vrouwtje**
Broedzorg	**Bewaakt de eieren, draagt de kikkervisjes en voedt de kikkervisjes met eieren**

VERZORGING

Min. terrarium	**60 × 60 × 60 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**22-24 °C**
Temperatuur ’s nachts	**17-19 °C**
Luchtvochtigheid	**70-100%**
Water	**Geen waterbak**
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Ernstig bedreigd (CR)**

Verzorging in het kort

HET TERRARIUM
A pair of *O. lehmanni* needs a vivarium of at least 60 × 60 × 60 cm (length × width × height), and larger is better, especially for breeding.

TEMPERATUUR
O. lehmanni comes from cool mountain rainforest and is more sensitive to heat than most poison frogs. It does best at 22-24 °C during the day, and 24-25 °C should not be exceeded, not even briefly.

LUCHTVOCHTIGHEID EN SPROEIEN
Humidity should be 70-100 %: high in the morning and evening, preferably with fog so that it reaches 100 %, and a little lower in the middle of the day, down to about 70 %.

WATER EN WATERKWALITEIT
The frogs do not need a water bowl, because the humidity covers their needs.

VERLICHTING EN UVB
The frog is diurnal and lives in shade under the forest canopy. There is no documentation that the species needs UVB when the food is dusted with vitamin D3, but weak UVB lighting (Ferguson zone 1) can be used without harm.

VOEDING
O. lehmanni eats only live, small arthropods. The staple food is flightless fruit flies, and the species is large enough to take the larger *Drosophila hydei* as well.

GEDRAG EN OMGANG
O. lehmanni is diurnal and calm, but the males are territorial and call from fixed spots. Keep the species in pairs.

NATUURLIJK VERSPREIDINGSGEBIED
Colombia.