herpy

PIJLGIFKIKKERS

Strawberry Poison Frog

Oophaga pumilio

Oophaga pumilio is a small, diurnal poison frog of 17-24 mm from the rainforests of Central America, known for its many colour forms and an unusual form of parental care: the female feeds each tadpole with unfertilised eggs. It is fascinating to watch, but requires experience, small feeder insects and patience.

Familie: Dendrobatidae Vroegere naam: *Dendrobates pumilio*

KERNGEGEVENS

HET DIER

Volwassen grootte	**17-24 mm** kop-romplengte
Levensduur	**10-15 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Geslachtsrijp	**10 maanden**
Eieren per legsel	**3-5**
Uitkomen	**10-14 dagen**
Larvenperiode	**42-56 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Onbevruchte eieren van het vrouwtje**
Broedzorg	**Bewaakt de eieren, draagt de kikkervisjes en voedt de kikkervisjes met eieren**

VERZORGING

Min. terrarium	**45 × 45 × 45 cm** L × B × H voor 4 volwassen dieren
Temperatuur overdag	**22-26 °C**
Temperatuur ‘s nachts	**19-21 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	A pair or a small group of up to four animals needs a vivarium of at least 45 × 45 × 45 cm (length × width × height). More animals require more floor area.
TEMPERATUUR	By day the temperature should be 22-26 °C, preferably about 24 °C, and at night 19-21 °C. Temperatures below 18 °C or above 32 °C for any length of time can be fatal.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be high, 70-100 %. Aim for 70-80 % during the day and up to 100 % at night, for example by misting four times a day with an automatic system. A timer-controlled rain system is therefore a great advantage.
WATER EN WATERKWALITEIT	A shallow water dish is enough, but the water must be no more than about ¾ of the frog's height, in practice under 1 cm, because the species swims poorly and can drown.
VERLICHTING EN UVB	Give 12 hours of light a day, so that the plants thrive. An LED plant lamp that does not give off much heat is a good choice.
VOEDING	*Oophaga pumilio* needs very small feeder insects, at most about 2 mm. Small, flightless *Drosophila melanogaster* and springtails are the staple, supplemented with aphids, termites and other small feeders.
GEDRAG EN OMGANG	*Oophaga pumilio* is diurnal, bold and inquisitive and moves about a great deal in the leaf litter. The males defend territories with a buzzing, cricket-like call and may fight other males.
NATUURLIJK VERSPREIDINGSGEBIED	Nicaragua, Costa Rica en Panama.