herpy

PIJLGIFKIKKERS

Little Devil Poison Frog

Oophaga sylvatica

Oophaga sylvatica is one of the largest poison frogs in the genus *Oophaga*: a diurnal forest-floor frog of 3-4 cm from the rainforests of Ecuador and Colombia, known for its many colour forms in orange, red and yellow on a black or brown background and for its persistent brood care.

Familie: Dendrobatidae Vroegere naam: *Dendrobates sylvaticus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**3-4 cm** kop-romplengte
Levensduur	**10-15 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Geslachtsrijp	**18-24 maanden**
Eieren per legsel	**4-20**
Uitkomen	**10-16 dagen**
Larvenperiode	**75-90 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Onbevruchte eieren van het vrouwtje**
Broedzorg	**Bewaakt de eieren, draagt de kikkervisjes en voedt de kikkervisjes met eieren**

VERZORGING

Min. terrarium	**60 × 60 × 60 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**23-26 °C**
Temperatuur ‘s nachts	**21-23 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Gevoelig (NT)**

Verzorging in het kort

HET TERRARIUM
An enclosure of at least 60 × 60 × 60 cm (length, width, height) suits one pair. The males are strongly territorial, and the females also hold territories and form a hierarchy when there are many animals.

TEMPERATUUR
Oophaga sylvatica comes from a lowland with almost constant warmth, so the temperature must be kept stable: 23-26 °C by day and 21-23 °C at night.

LUCHTVOCHTIGHEID EN SPROEIEN
Humidity is kept at 70-80 % during the day and rises towards 100 % early in the morning and late in the evening, as when fog and rain drift in over the forest. At midday it may fall to about 70 %.

WATER EN WATERKWALITEIT
A shallow water dish with 1-2 cm of water and a ramp of stones or moss is enough for adults; the frogs are poor swimmers, and the water should reach no more than about three quarters of the animal's height when it stands.

VERLICHTING EN UVB
The lighting should first and foremost suit the plants. Fluorescent tubes or LED lights of about 6,500 K give natural colours and good plant growth; choose light sources with high efficiency and a reflector.

VOEDING
Adults take the usual small feeder insects: springtails, small and large fruit flies, micro crickets, small waxworm larvae, booklice, firebrats and finely sifted meadow plankton.

GEDRAG EN OMGANG
The males are strongly territorial and call from morning to evening; the call is short and high, about five notes per second. Rival males call in competition, advance and retreat and can end up in wrestling matches.

NATUURLIJK VERSPREIDINGSGEBIED
Ecuador en Colombia.