herpy

PIJLGIFKIKKERS

Vicente's Poison Frog

Oophaga vicentei

Oophaga vicentei is a small, brightly coloured and diurnal poison dart frog of up to 24 mm, found only in central Panama. The colours vary from yellow and red to blue, grey and green, often with a black reticulated pattern. The species is endangered in the wild, and little is known about its husbandry.

Familie: Dendrobatidae Vroegere naam: *Dendrobates vicentei*

KERNGEGEVENS

HET DIER

Volwassen grootte	**12-24 mm** kop-romplengte
Levensduur	**10-15 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend en bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**In bladoksels (bijv. bromelia's)**
Eieren per legsel	**1-12**
Uitkomen	**10-14 dagen**
Larvenperiode	**42-56 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Onbevruchte eieren van het vrouwtje**
Broedzorg	**Bewaakt de eieren, draagt de kikkervisjes en voedt de kikkervisjes met eieren**

VERZORGING

Min. terrarium	**45 × 45 × 60 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**22-26 °C**
Temperatuur 's nachts	**19-21 °C**
Luchtvochtigheid	**80-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Bedreigd (EN)**

Verzorging in het kort

HET TERRARIUM	Little is known about the husbandry of *Oophaga vicentei*.
TEMPERATUUR	The daytime temperature should be 22-26 °C and the night 19-21 °C. The frogs tolerate heat poorly, so avoid longer periods above 28 °C.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 80-100 % and never below 70 %. Mist the enclosure thoroughly once or twice a day with water free of lime and chlorine (rainwater or reverse-osmosis water), so that leaves and bromeliads are wet.
WATER EN WATERKWALITEIT	A shallow water bowl with 1-2 cm of water is enough, because the frogs are poor swimmers. Keep it clean, and change the water every day or every other day.
VERLICHTING EN UVB	UVB is hardly necessary, but a little UVB (Ferguson zone 1) does no harm and is optional. Plant lighting for 10-12 hours a day keeps the plants alive, and the night must be dark. The lamps must not heat the enclosure above 26 °C.
VOEDING	The species needs very small feeder insects. The staple is small, flightless fruit flies (*Drosophila melanogaster*) and springtails, supplemented with aphids, small termites and small isopods.
GEDRAG EN OMGANG	*Oophaga vicentei* is diurnal. The male calls from a spot high up and finds the breeding site. Females may eat other females' eggs, so a pair is the safest. The frogs should not be handled, because the skin is thin and sensitive.
NATUURLIJK VERSPREIDINGSGEBIED	Panama.