herpy

PIJLGIFKIKKERS

Kokoe Poison Frog

Phyllobates aurotaenia

Phyllobates aurotaenia is a diurnal poison frog of 3-4 cm from the Pacific coast of Colombia, black with two green to golden stripes along the back. Wild animals carry a deadly skin poison, but captive-bred animals are non-poisonous. The species is bold, calls loudly and is easy to breed.

Familie: Dendrobatidae Vroegere naam: *Dendrobates aurotaenia*

KERNGEGEVENS

HET DIER

Volwassen grootte	**3-4 cm** kop-romplengte
Levensduur	**10-20 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Sterk**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Geslachtsrijp	**8-10 maanden**
Eieren per legsel	**8-28**
Larvenperiode	**60-80 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kant-en-klaar voer**
Broedzorg	**Draagt de kikkervisjes**

VERZORGING

Min. terrarium	**60 × 45 × 45 cm** L × B × H voor 4 volwassen dieren
Temperatuur overdag	**21-24 °C**
Temperatuur ’s nachts	**18-21 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

A pair or a small group of *Phyllobates aurotaenia* needs a rainforest enclosure of at least 60 × 45 × 45 cm for up to 4 adults. The species is active and also uses height, so give it perches and plants to climb in.

TEMPERATUUR

Phyllobates aurotaenia does best when cool. During the day it should be 21-24 °C, and at night the temperature is lowered to 18-21 °C. The temperature must not exceed 27 °C, because the genus tolerates heat poorly.

LUCHTVOCHTIGHEID EN SPROEIEN

Humidity should be 70-100 %. The species tolerates down to 50 % briefly if it has access to water, but low humidity without water can quickly become fatal.

WATER EN WATERKWALITEIT

A shallow water dish with 1-2 cm of water and a stone or a sloping edge, so the frogs can easily get up, is enough.

VERLICHTING EN UVB

The light is first and foremost for the plants. Fluorescent tubes or LED light of about 6,500 K give natural colours and good plant growth, and about 12 hours of light a day is usual.

VOEDING

Phyllobates aurotaenia eats small arthropods. Juveniles take wingless *Drosophila melanogaster*, while adults readily eat larger feeder insects such as *Drosophila hydei* and very small crickets.

GEDRAG EN OMGANG

Phyllobates aurotaenia is diurnal and bold, but shy when young.

NATUURLIJK VERSPREIDINGSGEBIED

Colombia.