herpy

PIJLGIFKIKKERS

Black-legged Poison Dart Frog

Phyllobates bicolor

Phyllobates bicolor is a yellow to orange poison frog of 3-5 cm from the rainforest of western Colombia, often with black or greenish legs. In the wild it is one of the most poisonous animals known, but captive-bred animals are non-poisonous. It is bold, curious and almost always on view, and it breeds readily in the enclosure.

Familie: Dendrobatidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**3-5 cm** kop-romplengte
Gewicht	**4,5 g**
Levensduur	**6-20 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Sterk**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Geslachtsrijp	**12 maanden**
Eieren per legsel	**4-30**
Uitkomen	**13-14 dagen** bij 21-23 °C
Larvenperiode	**21-60 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**50 × 50 × 50 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**21-25 °C**
Temperatuur ’s nachts	**18-21 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Insecten en kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Bedreigd (EN)**

Verzorging in het kort

HET TERRARIUM	A pair of *Phyllobates bicolor* needs a rainforest enclosure of at least 50 × 50 × 50 cm. Allow about 40 litres per adult. A group of up to nine animals needs at least 70 × 70 × 70 cm.
TEMPERATUUR	During the day it should be 21-25 °C, and at night the temperature is lowered by 3-4 degrees, to 18-21 °C.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 70-80 % in the middle of the day and up to 100 % morning and evening. A rain system and a fogging system on timers make this easiest, and the fog clearly makes the frogs call and move about.
WATER EN WATERKWALITEIT	The species appreciates a water surface in the enclosure.
VERLICHTING EN UVB	The lighting is first and foremost for the plants. Fluorescent tubes or LED for about 12 hours a day are enough.
VOEDING	*Phyllobates bicolor* prefers small feeder insects of no more than about 3 mm.
GEDRAG EN OMGANG	*Phyllobates bicolor* is diurnal, very active and curious and almost always on view. The males call loudly with trilling, buzzing tones, sometimes for several days at a time.
NATUURLIJK VERSPREIDINGSGEBIED	Colombia.