herpy

PIJLGIFKIKKERS

Golden Poison Frog

Phyllobates terribilis

Phyllobates terribilis is the largest of the poison frogs, 4-5 cm, and in the wild one of the most toxic animals known. It lives only in a small patch of rainforest on Colombia's Pacific coast. Captive-bred animals are practically non-toxic, bold and always on show, and the species has become one of the most commonly kept poison frogs in the hobby.

Familie: Dendrobatidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**4-5 cm** kop-romplengte
Levensduur	**8-15 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Sterk**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Geslachtsrijp	**16-20 maanden**
Eieren per legsel	**10-30**
Uitkomen	**10-25 dagen** bij 22-24 °C
Larvenperiode	**30-60 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**60 × 50 × 50 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**21-25 °C**
Temperatuur 's nachts	**18-21 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Insecten en kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Bedreigd (EN)**

Verzorging in het kort

HET TERRARIUM	A pair of *Phyllobates terribilis* needs an enclosure of at least 60 × 50 × 50 cm. A group of 4-6 animals needs at least 90 × 50 × 50 cm, and more animals require more floor area.
TEMPERATUUR	During the day it should be 21-25 °C, preferably a couple of degrees warmer at the top of the enclosure, and at night 18-21 °C, which corresponds to room temperature. The temperature must not exceed 27 °C for any length of time.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be high, 70-100 %, but the floor must not become boggy. A rain system is the easiest, for example with three mistings a day of 20-60 seconds, depending on the nozzles.
WATER EN WATERKWALITEIT	Provide one or more small water dishes with about 2 cm of water, for example saucers or halved Brazil nut shells, which the animals can easily climb out of.
VERLICHTING EN UVB	The light is first and foremost for the plants. Fluorescent tubes or LED bars for about 12 hours a day are enough, and the day length may vary slightly over the year.
VOEDING	*Phyllobates terribilis* takes larger prey than most poison dart frogs. In the wild ants are the main food, but it takes many kinds of small arthropods.
GEDRAG EN OMGANG	*Phyllobates terribilis* is diurnal, bold and inquisitive and is out on show most of the day with a short break in the middle of the day. The males call with a soft, bird-like trill, especially after misting.
NATUURLIJK VERSPREIDINGSGEBIED	Colombia.