herpy

PIJLGIFKIKKERS

Golfodulcean Poison Frog

Phyllobates vittatus

Phyllobates vittatus is a black poison frog with two bright orange or yellow stripes down the back and blue marbled legs. It occurs only in the lowland forest around the Golfo Dulce in south-western Costa Rica. In the terrarium it is diurnal, vocal and a prolific breeder, and the male carries the tadpoles to water himself.

Familie: Dendrobatidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**3-4 cm** kop-romplengte
Levensduur	**10-20 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Geslachtsrijp	**8-10 maanden**
Eieren per legsel	**7-21**
Larvenperiode	**45-80 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**60 × 45 × 45 cm** L × B × H voor 4 volwassen dieren
Temperatuur overdag	**21-25 °C**
Temperatuur ’s nachts	**18-21 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Insecten en kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Kwetsbaar (VU)**

Verzorging in het kort

HET TERRARIUM	Up to four *Phyllobates vittatus* need an enclosure of at least 60 × 45 × 45 cm. A group of 5-6 animals needs more space, e.g. 100 × 60 × 60 cm, which also gives room both to move on the ground and to climb.
TEMPERATUUR	During the day *Phyllobates vittatus* needs 21-25 °C and at night 18-21 °C, which most living rooms provide.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 70-100 %. The species tolerates down to 50 % for short periods if it has access to water, but low humidity without water can quickly become fatal.
WATER EN WATERKWALITEIT	Provide a shallow water dish or a small water feature with 1-2 cm of water and a stone or a piece of bark so the animals can climb out easily. The frogs moisten themselves in the water, and the male may use it to deposit tadpoles.
VERLICHTING EN UVB	The lighting is for the plants. LED bars or fluorescent tubes on a timer for 10-12 hours a day are enough, and the night must be dark. Provide shade, and never place the enclosure in direct sun.
VOEDING	*Phyllobates vittatus* is unusual in that adults take larger prey than most poison frogs.
GEDRAG EN OMGANG	*Phyllobates vittatus* is diurnal, and the males have a high, trilling call. The species can be kept in pairs or groups.
NATUURLIJK VERSPREIDINGSGEBIED	Costa Rica.