herpy

PIJLGIFKIKKERS

Amazonian Poison Frog

Ranitomeya amazonica

Ranitomeya amazonica is a tiny, diurnal poison frog of 16-19 mm from the Amazon around Iquitos in Peru and Leticia in Colombia. It is black with red, orange or yellow stripes and blue-grey marbling on the legs, breeds in bromeliads and is regarded as one of the easiest small Ranitomeya to keep and breed.

Familie: Dendrobatidae Vroegere naam: *Dendrobates amazonicus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**16-19 mm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend en bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**In bladoksels (bijv. bromelia's)**
Geslachtsrijp	**5-8 maanden**
Eieren per legsel	**2-8**
Uitkomen	**10-14 dagen**
Larvenperiode	**60-80 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kant-en-klaar voer en andere kikkervisjes**
Broedzorg	**Draagt de kikkervisjes**

VERZORGING

Min. terrarium	**40 × 40 × 40 cm** L × B × H voor 3 volwassen dieren
Temperatuur overdag	**21-26 °C**
Temperatuur 's nachts	**18-21 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel**
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

A pair or trio of *Ranitomeya amazonica* needs an enclosure of at least 40 × 40 × 40 cm. If there are to be more frogs, allow about 20 litres of extra space per animal, and choose a tall rather than a low enclosure.

TEMPERATUUR

Keep the enclosure at 21-26 °C by day and 18-21 °C at night. The frogs are most active and breed best at the warm end, at 24-26 °C.

LUCHTVOCHTIGHEID EN SPROEIEN

Humidity should be 70-100 %, preferably 80-100 %. Mist every day, preferably morning and evening, so that the humidity is close to 100 % after misting and falls to about 70-75 % in the middle of the day.

WATER EN WATERKWALITEIT

Ranitomeya amazonica swims poorly, so the water must be shallow and easy to climb out of.

VERLICHTING EN UVB

The frogs are diurnal and live in the shade of the forest. The lighting is first and foremost for the plants: give 12-14 hours of LED plant lighting a day. *Ranitomeya amazonica* has no known need for UVB, so UVB is optional.

VOEDING

Ranitomeya amazonica is so small that it can only take small feeder insects. Juveniles and adults take wingless fruit flies (*Drosophila melanogaster*), and large adults can also take *D. hydei* and newly hatched crickets.

GEDRAG EN OMGANG

Ranitomeya amazonica is diurnal, but often shy, and young animals in particular hide a lot. In a densely planted enclosure with many dead leaves the frogs come out more.

NATUURLIJK VERSPREIDINGSGEBIED

Peru en Colombia.