herpy

PIJLGIFKIKKERS

Tahuayo Poison Frog

Ranitomeya flavovittata

Ranitomeya flavovittata is one of the smallest species in the genus: only 15-17 mm from snout to vent. It is bright yellow with black markings, brown legs and a pale blue belly and comes from the rainforest near Iquitos in north-eastern Peru. It is diurnal and a little shy, breeds willingly in film canisters and bromeliads and is a good choice for a keeper with a little experience.

Familie: Dendrobatidae Vroegere naam: *Dendrobates flavovittatus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**15-17 mm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend en bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**In bladoksels (bijv. bromelia's)**
Geslachtsrijp	**5-8 maanden**
Eieren per legsel	**2-6**
Uitkomen	**10-14 dagen**
Larvenperiode	**60-80 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kant-en-klaar voer en onbevruchte eieren van het vrouwtje**
Broedzorg	**Draagt de kikkervisjes en voedt de kikkervisjes met eieren**

VERZORGING

Min. terrarium	**30 × 30 × 45 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**21-26 °C**
Temperatuur 's nachts	**18-21 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel**
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

A pair of *R. flavovittata* needs an enclosure of at least 30 × 30 × 45 cm, but bigger is always better. A group of 3-4 frogs needs at least 45 × 45 × 45 cm, and each additional frog requires more space.

TEMPERATUUR

Keep the enclosure at 21-26 °C by day and 18-21 °C at night. The night temperature is borrowed from related species. The temperature must not exceed 27 °C for long periods, and above 29 °C can be fatal.

LUCHTVOCHTIGHEID EN SPROEIEN

Humidity must be 70-100 %. The species can briefly cope with about 50 % if it has access to clean water, but this is an emergency measure and can be fatal if it continues. Mist daily, and use reverse-osmosis water or rainwater.

WATER EN WATERKWALITEIT

Give the frogs a shallow water dish or a small water feature, and keep the water so shallow that the smallest frog can sit with its head and upper body above the water: the species swims poorly. The water depth should be 1-2 cm.

VERLICHTING EN UVB

The frogs are diurnal and live in the shade of the rainforest. The light is first and foremost for the plants: LED light for 12-14 hours a day is suitable.

VOEDING

R. flavovittata is so small that it can only take small feeder insects. Juveniles and most adults eat wingless fruit flies (*Drosophila melanogaster*), and large adults can also take *D. hydei* and newly hatched crickets.

GEDRAG EN OMGANG

R. flavovittata is diurnal, quick and a little shy. The frogs become bolder if the enclosure is densely planted and has many hiding places, and they get used to you if you watch them through the glass.

NATUURLIJK VERSPREIDINGSGEBIED

Peru.