herpy

PIJLGIFKIKKERS

Red-backed Poison Frog

Ranitomeya reticulata

Ranitomeya reticulata is one of the smallest poison frogs: only 13-17 mm, with a coppery red to orange back and blue-grey legs with a black net pattern. It lives in the lowland forest of the Amazon in north-eastern Peru and eastern Ecuador. In the terrarium it is bold and lively, but demanding, because the eggs are hard to find and the juveniles are hard to raise.

Familie: Dendrobatidae Vroegere naam: *Dendrobates reticulatus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**13-17 mm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend en bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**1-8**
Uitkomen	**10-14 dagen**
Larvenperiode	**60-80 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Draagt de kikkervisjes**

VERZORGING

Min. terrarium	**30 × 30 × 45 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**22-26 °C**
Temperatuur ’s nachts	**18-22 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel**
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	A pair of *R. reticulata* needs an enclosure of at least 30 × 30 × 45 cm, preferably 40 × 40 × 45 cm or larger. The frogs like to climb, so the height is used.
TEMPERATUUR	Keep the enclosure at 22-26 °C during the day and 18-22 °C at night. The temperature must not exceed 27 °C for any length of time, and above 29 °C can be fatal. Heat is always more dangerous than cold.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 70-100 %. A good rhythm is almost 100 % morning and evening and 70-75 % in the middle of the day, so the enclosure dries out a little between sprayings.
WATER EN WATERKWALITEIT	A shallow water dish or a bromeliad with water in the leaf axils is enough.
VERLICHTING EN UVB	*R. reticulata* is diurnal, and the lighting is first and foremost for the plants. Give 12-14 hours of light a day, e.g. LED from 8 to 20 hrs, and darkness at night. It is not documented that the species needs UVB.
VOEDING	The staple diet is small arthropods. Juveniles and most adults take flightless fruit flies (*Drosophila melanogaster*), and large adults can also take the larger *D. hydei*.
GEDRAG EN OMGANG	*R. reticulata* is lightning fast and bold and is often seen out in the open, as if it has no idea how small it is. Males call from fixed spots with a high, short buzz.
NATUURLIJK VERSPREIDINGSGEBIED	Peru en Ecuador.

De volledige gids met kaart en updates: herpy.org/en/poison-dart-frogs/ranitomeya-reticulata