herpy

PIJLGIFKIKKERS

Sira Poison Frog

Ranitomeya sirensis

Ranitomeya sirensis is a small, diurnal poison frog of 15-25 mm from eastern Peru, northern Bolivia and the far west of Brazil. The species includes the forms that were formerly sold as "lamasi" and "biolat", and most of them are among the easiest poison frogs to breed. The frogs thrive in groups when there is enough space.

Familie: Dendrobatidae Vroegere naam: *Ranitomeya lamasi; Dendrobates lamasi; Dendrobates sirensis*

KERNGEGEVENS

HET DIER

Volwassen grootte	**15-25 mm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend en bodembewonend**
Leefgebied	**Regenwoud en bergbos**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**In bladoksels (bijv. bromelia's)**
Geslachtsrijp	**5-8 maanden**
Eieren per legsel	**1-8**
Uitkomen	**10-14 dagen**
Larvenperiode	**58-80 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**45 × 45 × 45 cm** L × B × H voor 3 volwassen dieren
Temperatuur overdag	**21-25 °C**
Temperatuur 's nachts	**18-21 °C**
Luchtvochtigheid	**80-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel**
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

A pair or trio of *R. sirensis* needs an enclosure of at least 45 × 45 × 45 cm. The species benefits from a little extra floor space and dense plant cover, and the large highland form thrives best in a larger enclosure.

TEMPERATUUR

Keep the enclosure at 21-25 °C during the day and 18-21 °C at night. The temperature must not exceed 27 °C for any length of time, and above 29 °C is dangerous.

LUCHTVOCHTIGHEID EN SPROEIEN

Humidity should be 80-100 %. The species tolerates down to 50 % only briefly, and only if it has access to water; low humidity without water can quickly become fatal.

WATER EN WATERKWALITEIT

A shallow water dish or a bromeliad with water in the leaf axils is enough. Keep the water 1-2 cm deep, so the smallest frog can sit in it with its head and upper body above the water.

VERLICHTING EN UVB

R. sirensis is diurnal, and the lighting is first and foremost for the plants. Give 12-14 hours of light a day with fluorescent tubes or LED and darkness at night. It is not documented that the species needs UVB.

VOEDING

The staple diet is small arthropods: wingless fruit flies (*Drosophila melanogaster*), springtails and small woodlice. Adults also take the larger *D. hydei* and newly hatched crickets.

GEDRAG EN OMGANG

R. sirensis is often shy and likes to hide, so a densely planted enclosure with many hiding places gives the most to look at. The male calls to lure a female to an egg-laying site.

NATUURLIJK VERSPREIDINGSGEBIED

Bolivia, Brazilië en Peru.