herpy

PIJLGIFKIKKERS

Summers' Poison Dart Frog

Ranitomeya summersi

Ranitomeya summersi is a small, diurnal poison frog of 16-21 mm from the Huallaga Valley in Peru, orange or copper-coloured with black marbling and a black mask over the eyes. The species is threatened in the wild, the eggs are hard to find, and the frogs are quick and shy, so it is best suited to a keeper with experience of poison frogs.

Familie: Dendrobatidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**16-21 mm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend en bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**4-9**
Uitkomen	**10-14 dagen**
Larvenperiode	**60-80 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**30 × 30 × 45 cm** L × B × H voor 3 volwassen dieren
Temperatuur overdag	**21-26 °C**
Temperatuur 's nachts	**18-21 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel**
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Bedreigd (EN)**

Verzorging in het kort

HET TERRARIUM	A pair or a small trio of *R. summersi* can be kept in an enclosure of at least 30 × 30 × 45 cm (length × width × height), and bigger is better.
TEMPERATUUR	Keep the enclosure at 21-26 °C during the day and 18-21 °C at night. The temperature must not exceed 27 °C for any length of time, and above 29 °C is life-threatening. The species tolerates heat worse than cold.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 70-100 %. A good rhythm is close to 100 % morning and evening and 70-75 % in the middle of the day, so the enclosure can dry out a little between mistings.
WATER EN WATERKWALITEIT	A shallow water dish or a bromeliad with water in the leaf axils is enough.
VERLICHTING EN UVB	*R. summersi* is diurnal, and the light is first and foremost for the plants. Give 12-14 hours of light a day and darkness at night. It is not documented that the species needs UVB.
VOEDING	The staple food is small live arthropods. Juveniles and young animals take wingless fruit flies (*Drosophila melanogaster*), and large adults can also take the larger *D. hydei*.
GEDRAG EN OMGANG	*R. summersi* is diurnal, quick and often shy, but can become bolder once it has got used to the enclosure. They are seen most in a densely planted enclosure with hiding places.
NATUURLIJK VERSPREIDINGSGEBIED	Peru.