herpy

PIJLGIFKIKKERS

Red Uakari Poison Frog

Ranitomeya uakarii

Ranitomeya uakarii is a very small, diurnal poison frog of 13-16 mm from the Amazon basin. It is black with red to orange stripes, yellow flanks and blue legs, and it is rare in the hobby. The care advice is based on experience with the genus, and the species suits an experienced keeper with patience and a steady supply of tiny feeder insects.

Familie: Dendrobatidae Vroegere naam: *Dendrobates uakarii*

KERNGEGEVENS

HET DIER

Volwassen grootte	**13-16 mm** kop-romplengte
Levensduur	**5-6 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend en bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**In bladoksels (bijv. bromelia's)**
Eieren per legsel	**2-7**
Uitkomen	**10-14 dagen**
Larvenperiode	**60-100 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Draagt de kikkervisjes**

VERZORGING

Min. terrarium	**45 × 45 × 45 cm** L × B × H voor 3 volwassen dieren
Temperatuur overdag	**21-25 °C**
Temperatuur 's nachts	**18-21 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel**
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM
A pair or trio of *Ranitomeya uakarii* needs an enclosure of at least 45 × 45 × 45 cm. The size is borrowed from the genus. If you use juveniles, the safest approach is to start with at least four, so that both sexes are included.

TEMPERATUUR
Keep the enclosure at 21-25 °C during the day and 18-21 °C at night. The figures are borrowed from the genus. Extra heat is rarely necessary in a heated living room.

LUCHTVOCHTIGHEID EN SPROEIEN
Humidity should be 70-100 %. Mist daily, so that humidity is close to 100 % morning and evening and 70-75 % in the middle of the day. This lets the terrarium dry out a little between sprayings.

WATER EN WATERKWALITEIT
A shallow water dish or a bromeliad with water in the leaf axils is enough. Keep the water 1-2 cm deep, so that the small frog can sit in it with its head and upper body above the water; the depth is borrowed from the genus.

VERLICHTING EN UVB
Ranitomeya uakarii is diurnal, and the lighting is first and foremost for the plants. Provide 12-14 hours of light a day with fluorescent tubes or LED and darkness at night. It is not documented that the species needs UVB.

VOEDING
The staple diet is small arthropods: wingless fruit flies, springtails, small white woodlice and newly hatched crickets.

GEDRAG EN OMGANG
Ranitomeya uakarii is diurnal. In a densely planted terrarium with many hiding places you typically see more of the frogs than in a barren terrarium, but the small frogs are often shy and hide when you approach the glass.

NATUURLIJK VERSPREIDINGSGEBIED
Brazilië, Colombia en Peru.

De volledige gids met kaart en updates: herpy.org/en/poison-dart-frogs/ranitomeya-uakarii