herpy

SALAMANDERS EN WATERSALAMANDERS

Guangxi Warty Newt

Paramesotriton guangxiensis

Paramesotriton guangxiensis is a robust, almost entirely aquatic salamander of 13-15 cm from forest streams and rivers in Guangxi in southern China. It is dark brown with granular skin, rare and known from only a few localities. Little is known about its care, but the species is kept like the other species in the genus: in cool, clean water below 22 °C.

Familie: Salamandridae Vroegere naam: *Trituroides guangxiensis* **Mild huidgif**

KERNGEGEVENS

HET DIER

Volwassen grootte	**13-15 cm** totale lengte incl. staart
Levensduur	**10-15 jaar**
Actief	**In de schemering en 's nachts**
Levenswijze	**Aquatisch**
Leefgebied	**Regenwoud en moerasgebied**
Huidgif	**Mild** Was je handen na contact en houd je vingers uit de buurt van je ogen en mond
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**60 × 30 × 35 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**15-20 °C**
Temperatuur 's nachts	**13-18 °C**
Luchtvochtigheid	**70-90%**
Water	**Aquaterrarium** diepte 15-20 cm
UVB	**Niet nodig**
Voeding	**Insecten en wormen/slakken**
Gehouden	**In paren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Larven in het water**
Legplaats	**In het water**
Eieren per seizoen	**20-115**
Legsels per jaar	**1**
Uitkomen	**21-56 dagen** bij 18-22 °C
Larvenperiode	**80-110 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kleine waterongewervelden (watervlooien, muggenlarven)**
Broedzorg	**Geen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
EU-bijlage	**Bijlage B**
IUCN	**Kwetsbaar (VU)**

Over de soort

Paramesotriton guangxiensis is a medium-sized, sturdily built salamander. Females reach about 13 cm in total length, males a little more. The skin is dark brown, the head broad, and the tail tall and laterally compressed, as in the rest of the genus. The animals live in the water almost their whole lives.

The skin secretes a toxic substance. Tetrodotoxin has been detected in the skin of several species in the genus, but the toxin has not been studied in this species. Wash hands after contact, do not touch the eyes or mouth, and handle the animals only when necessary.

The species was described in 1983 by Huang, Tang and Tang. It is rare, threatened by habitat loss and very little known in captivity. Genetic and morphological studies suggest that the populations in Quang Ninh in north-eastern Vietnam, previously assigned to *Paramesotriton deloustali*, may belong to this species. Lifespan is not documented; the figure in the box is borrowed from the genus, where 10-15 years is common.

Verspreiding en leefgebied

Paramesotriton guangxiensis is reliably known only from Paiyangshan in Ningming in Guangxi in southern China, close to the border with Vietnam. The populations in Quang Ninh in north-eastern Vietnam may also belong to the species, but this has not been resolved, and the map shows only China.

> **Natuurlijk verspreidingsgebied:** China.

LEEFGEBIED

Paramesotriton guangxiensis lives in streams and rivers in humid subtropical to tropical lowland forest. The animals stay in clean, cool, oxygen-rich water with stones and hides. The climate is seasonal, with hot, rainy summers and cooler winters. Like the rest of the genus, the species probably tolerates heat poorly. The habitats are threatened by deforestation and agriculture.

Verzorging

HET TERRARIUM

Little is known about the care of *Paramesotriton guangxiensis*. The species is rare in captivity, and the figures here are borrowed from the closely related *Paramesotriton chinensis*, *Paramesotriton deloustali* and *Paramesotriton hongkongensis*, which are far better known and have the same requirements for cool, clean water.

A pair needs an aquarium of at least 60 × 30 × 35 cm (about 60 litres), and more animals require more space. Provide plenty of hides of stone, bark and roots, and a small land area of a stone or a floating cork island. Keep the lid tight, because the animals can leave the water. Use sand, large stones or no substrate, because fine gravel can be swallowed and cause intestinal blockage.

TEMPERATUUR

Paramesotriton guangxiensis must be kept cool: 15-20 °C by day and 13-18 °C at night, and the water must not become warmer than 22 °C. In summer, keeping the aquarium below the limit may require a fan, cooling elements or a cool room. True brumation is not documented for the species.

LUCHTVOCHTIGHEID EN SPROEIEN

Paramesotriton guangxiensis is almost entirely aquatic, so the humidity of the room plays a small role; 70-90 % is fine. Any land area must be damp.

WATER EN WATERKWALITEIT

The water should be 15-20 cm deep. Use dechlorinated tap water. The water must be clean and oxygen-rich, without extreme pH or hardness values. A sponge filter driven by an air pump or a small internal filter with a gentle current suits the species. Change part of the water regularly, and remove uneaten food.

VERLICHTING EN UVB

Paramesotriton guangxiensis does not need strong light. A single fluorescent or LED tube on a timer provides a fixed daily rhythm of about 12 hours. UVB is not necessary.

Lees meer: Licht en UVB

VOEDING

Paramesotriton guangxiensis eats live or chopped earthworms, crickets, freshly moulted mealworms, waxworms and pieces of shrimp, like the rest of the genus. Adults are fed twice a week, juveniles three to four times a week. Accustom the animals to taking food from tweezers, so that it is easy to see how much each animal eats. The animals easily become too fat, so do not feed waxworms alone.

GEDRAG EN OMGANG

The behaviour of *Paramesotriton guangxiensis* is not described; species in the genus are often shy at first, but eat readily and quickly become confident. Keep males apart if they show aggression. Do not handle the animals more than necessary, and use a net when they have to be moved.

Kweek

GESLACHTSRIJPHEID EN GESLACHTSONDERSCHEID

In *Paramesotriton guangxiensis* the female's tail is probably shorter than the male's, and in the breeding season the male develops a swollen vent and a coloured tail stripe, as in the rest of the genus. The age at sexual maturity is not documented; in closely related species it takes several years.

PARING

The breeding season and mating are not documented for the species. The pattern in the genus is that the male fans his tail in front of the female and deposits a spermatophore, which she picks up with her vent. The female must be able to withdraw from the male, so the aquarium must be large with many hides.

EIEREN EN LARVEN

The female of *Paramesotriton guangxiensis* probably lays the eggs singly on aquatic plants and folds the leaf around them, as is typical for the genus. The number of eggs is not documented for the species; the figure in the box is borrowed from *Paramesotriton hongkongensis*. Move the eggs to a tub with an air stone, or place them in a breeding net, and change the water regularly.

METAMORFOSE

Hatching time and larval period are not documented for the species. The figures in the box are borrowed from *Paramesotriton deloustali*: the eggs hatch after 3-8 weeks at 18-22 °C, and metamorphosis takes place after about 80-110 days. Keep eggs and larvae below 22 °C. The larvae are predators and take daphnia and mosquito larvae, and they can eat each other, so sort them by size. At metamorphosis the gills shrink and the animal moves onto land.

OPFOK VAN DE JONGEN

Newly metamorphosed young move onto land and need a box with damp kitchen paper and moss. They take small earthworms, newly hatched crickets and small insects. When the young have grown, move them to a shallow aquarium with stones that protrude above the water, and tempt them into the water with earthworms at the water's edge.

Gezondheid en veelgemaakte fouten

Typical mistakes and diseases in *Paramesotriton guangxiensis*:

- Heat and poor water: keep the temperature below 22 °C and the water clean.
- Fine gravel that is swallowed and causes intestinal blockage.
- Bsal (*Batrachochytrium salamandrivorans*): Asian salamanders can carry the fungus without symptoms, and it can be fatal to other caudates. New animals must be quarantined, and equipment and hands must be cleaned between aquariums.
- Animals that will not eat, or have skin sores or swellings, must see a vet with expertise in amphibians.

Lees meer: Quarantaine en hygiëne

Vraag altijd raad aan iemand die er verstand van heeft

Deze gids is een goed beginpunt, maar kan ervaring niet vervangen. Dieren verschillen, en veel van wat er echt toe doet, leer je van mensen die de soort al jaren houden. Praat met een kweker, een ervaren houder of een dierenarts die reptielen en amfibieën kent voordat je het dier haalt - en opnieuw als er iets niet pluis lijkt.