Rough-skinned Newt

Taricha granulosa

Taricha granulosa is a 13-22 cm long newt from the Pacific coast of North America, with a dark, granular back and a bright orange belly. The skin contains tetrodotoxin, one of the most potent toxins found in amphibians, and the animal is deadly poisonous if swallowed. The species alternates between water and land, can be kept in water all year round and must be kept cool.

Familie: Salamandridae Vroegere naam: *Salamandra granulosa; Triturus granulosus; Triturus similans* **Sterk huidgif**

KERNGEGEVENS

HET DIER

Volwassen grootte	**13-22 cm** totale lengte incl. staart
Levensduur	**12-18 jaar**
Actief	**Overdag en 's nachts**
Levenswijze	**Semi-aquatisch**
Leefgebied	**Moerasgebied en gematigd bos en grasland**
Huidgif	**Sterk** Vermijd huidcontact en was je handen grondig na werk in het terrarium
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Larven in het water**
Legplaats	**In het water**
Geslachtsrijp	**48-60 maanden**
Eieren per seizoen	**7-30**
Uitkomen	**21-42 dagen**
Larvenperiode	**60-150 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kleine waterongewervelden (watervlooien, muggenlarven)**
Broedzorg	**Geen**

VERZORGING

Min. terrarium	**60 × 30 × 40 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**18-21 °C**
Temperatuur 's nachts	**14-18 °C**
Luchtvochtigheid	**80-95%**
Water	**Aquaterrarium** diepte 20-30 cm
UVB	**Niet nodig**
Voeding	**Insecten, kant-en-klaar voer en wormen/slakken**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

Taricha granulosa is the most aquatic species in the genus. The animals alternate between land and water, some for months or years at a time, but from about one year of age they can be kept in water all year round.

TEMPERATUUR

Taricha granulosa loves cool conditions. In summer, water and air are kept at 18-21 °C. If it gets warmer, towards 24 °C, the animals move up onto land; prolonged periods above 21 °C should be avoided.

LUCHTVOCHTIGHEID EN SPROEIEN

The land section and the hides above the water must be damp, but not soaking. Humidity has not been measured for the species; an aquaterrarium with a lid and damp hides gives around 80-95 %.

WATER EN WATERKWALITEIT

The water must be 20-30 cm deep, and deeper water does no harm: it is almost impossible to give the species water that is too deep in an ordinary tank.

VERLICHTING EN UVB

The species does not need UVB. Ordinary daylight in the room, or a weak fluorescent tube by day and darkness at night, is enough. The tank must not stand in direct sun, and heat lamps must be avoided.

VOEDING

Taricha granulosa is an opportunistic predator. In the wild it eats amphipods, insects, snails, leeches, tadpoles and amphibian and fish eggs.

GEDRAG EN OMGANG

Activity varies: some animals are out by day, while others hide under roots and stones and hunt at night. In the breeding season the male holds on to the female in amplexus, which can last from hours to several days.

NATUURLIJK VERSPREIDINGSGEBIED

Verenigde Staten en Canada.