California Newt

Taricha torosa

Taricha torosa is a sturdy newt, 13-20 cm long, from the California coast, with a granular brown back and a yellow to orange belly. The skin contains tetrodotoxin, a nerve toxin with no antidote, and the animal can be deadly if swallowed. The species lives mostly on land and enters the water only to breed, and it must be kept cool.

Familie: Salamandridae Vroegere naam: *Cynops torosus; Notophthalmus torosus* **Sterk huidgif**

KERNGEGEVENS

HET DIER

Volwassen grootte	**13-20 cm** totale lengte incl. staart
Levensduur	**12-22 jaar**
Actief	**Overdag en 's nachts**
Levenswijze	**Semi-aquatisch**
Leefgebied	**Moerasgebied en gematigd bos en grasland**
Huidgif	**Sterk** Vermijd huidcontact en was je handen grondig na werk in het terrarium
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Larven in het water**
Legplaats	**In het water**
Geslachtsrijp	**36 maanden**
Eieren per seizoen	**7-30**
Uitkomen	**14-21 dagen**
Larvenperiode	**60-90 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kleine waterongewervelden (watervlooien, muggenlarven)**
Broedzorg	**Geen**

VERZORGING

Min. terrarium	**60 × 30 × 30 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**16-21 °C**
Temperatuur 's nachts	**14-18 °C**
Luchtvochtigheid	**70-90%**
Water	**Waterbak**
UVB	**Niet nodig**
Voeding	**Insecten en wormen/slakken**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Niet opgenomen**
IUCN	**Gevoelig (NT)**

Verzorging in het kort

HET TERRARIUM

There is relatively little knowledge about the care of *Taricha torosa*, so the winter temperatures are borrowed from the closely related *Taricha granulosa*, which is kept more often.

TEMPERATUUR

Taricha torosa likes the cool. By day the temperature should be 16-21 °C and at night 14-18 °C. The temperature must not exceed 24 °C, and longer periods above 21 °C must be avoided.

LUCHTVOCHTIGHEID EN SPROEIEN

The substrate should be damp, but the enclosure should not drip. Humidity is about 70-90%, and daily misting with dechlorinated water keeps the substrate moist.

WATER EN WATERKWALITEIT

A shallow dish of clean water must always be available, so that the animal can cover itself completely with water. Use tap water treated for chlorine and chloramine; distilled and demineralised water are not used.

VERLICHTING EN UVB

The species does not need UVB. Daylight in the room or a weak fluorescent tube during the day and darkness at night is enough.

VOEDING

Taricha torosa is an opportunistic predator that takes almost anything it can overpower. Earthworms are a complete diet in themselves, and compost worms (*Eisenia fetida*) are also taken readily.

GEDRAG EN OMGANG

Taricha torosa is calm, sociable and not aggressive towards others of its kind of the same size.

NATUURLIJK VERSPREIDINGSGEBIED

Verenigde Staten.