herpy

SALAMANDERS EN WATERSALAMANDERS

Black Knobby Newt

Tylototriton asperrimus

Tylototriton asperrimus is a dark, warty crocodile newt of 12-15 cm with knobbly rib protuberances and heavy dorsal ridges. The species lives in humid mountain regions in the Chinese province of Guangxi and has given its name to a whole group of closely related crocodile newts, which have only recently been split off as separate species.

Familie: Salamandridae Vroegere naam: *Echinotriton asperrimus; Pleurodeles asperrimus* **Mild huidgif**

KERNGEGEVENS

HET DIER

Volwassen grootte	**12-15 cm** totale lengte incl. staart
Levensduur	**15-25 jaar**
Actief	**In de schemering en 's nachts**
Levenswijze	**Bodembewonend**
Leefgebied	**Bergbos en moerasgebied**
Huidgif	**Mild** Was je handen na contact en houd je vingers uit de buurt van je ogen en mond
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Larven in het water**
Legplaats	**Op de grond**
Eieren per seizoen	**30-80**
Uitkomen	**20-36 dagen** bij 22-24 °C
Larvenperiode	**50-110 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Andere larven en kleine waterongewervelden (watervlooien, muggenlarven)**
Broedzorg	**Geen**

VERZORGING

Min. terrarium	**60 × 40 × 40 cm** L × B × H voor 3 volwassen dieren
Temperatuur overdag	**20-24 °C**
Temperatuur 's nachts	**18-21 °C**
Luchtvochtigheid	**80-100%**
Water	**Watergedeelte** diepte 1-5 cm
UVB	**Optioneel**
Winterrust	**8-12 °C** 8-12 weken
Voeding	**Insecten en wormen/slakken**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
EU-bijlage	**Bijlage B**
IUCN	**Gevoelig (NT)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Tylototriton asperrimus*.
TEMPERATUUR	In summer the enclosure is kept at 20-24 °C by day and 18-21 °C at night, which corresponds to summer in the range.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity is 80-100 %. The enclosure is misted daily with a plant mister, and there must be both damper and drier zones so that the air does not stand still.
WATER EN WATERKWALITEIT	The water section must have soft, slightly acidic water; rainwater or reverse-osmosis water suits. The pH of the water is not documented for *Tylototriton asperrimus*; *Tylototriton ziegleri* is kept at pH 6.4-6.8.
VERLICHTING EN UVB	Light is of no particular importance to the animals and is arranged for the plants: LED strips or fluorescent tubes for 12 hours in summer and 10-11 hours in winter. UVB is not necessary for this secretive species.
VOEDING	Adults are given compost worms, crickets, house crickets, waxworms, earthworms and isopods of suitable size about twice a week. In winter they eat irregularly and much less.
GEDRAG EN OMGANG	In the closely related species the animals live very secretively and are seen mostly when fed or when the males are in the water. Pairs or small groups can be kept together in a spacious enclosure with many hiding places.
NATUURLIJK VERSPREIDINGSGEBIED	China.