Guizhou Crocodile Newt

Tylototriton kweichowensis

Tylototriton kweichowensis is a robust, blackish-brown crocodile newt of 15-21 cm with three broad orange to reddish stripes along the body. It lives on land in the mountains of Guizhou and Yunnan and enters the water mainly to breed. It is best kept in a large enclosure with both land and water.

Familie: Salamandridae Vroegere naam: *Pleurodeles kweichowensis* **Mild huidgif**

KERNGEGEVENS

HET DIER

Volwassen grootte	**15-21 cm** totale lengte incl. staart
Levensduur	**10-20 jaar**
Actief	**In de schemering en 's nachts**
Levenswijze	**Bodembewonend en semi-aquatisch**
Leefgebied	**Bergbos en moerasgebied**
Huidgif	**Mild** Was je handen na contact en houd je vingers uit de buurt van je ogen en mond
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Larven in het water**
Legplaats	**Op bladeren boven het water**
Eieren per seizoen	**50-150**
Legsels per jaar	**1**
Uitkomen	**10-12 dagen** bij 25 °C
Larvenperiode	**80-100 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kleine waterongewervelden (watervlooien, muggenlarven)**
Broedzorg	**Geen**

VERZORGING

Min. terrarium	**90 × 45 × 45 cm** L × B × H voor 3 volwassen dieren
Temperatuur overdag	**18-25 °C**
Temperatuur 's nachts	**15-20 °C**
Luchtvochtigheid	**70-90%**
Water	**Aquaterrarium** diepte 10-15 cm
UVB	**Niet nodig**
Voeding	**Insecten en wormen/slakken**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
EU-bijlage	**Bijlage B**
IUCN	**Kwetsbaar (VU)**

Verzorging in het kort

HET TERRARIUM	*Tylototriton kweichowensis* needs a large enclosure, because the animals are big and heavy, need a solid land area with a hiding place for each animal, and need enough water for both larvae and a filter.
TEMPERATUUR	The species thrives at ordinary room temperature: 18-25 °C by day and 15-20 °C at night. In summer the day may well be at the high end, and in winter the animals are kept at 15-20 °C around the clock.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity must be high, but the substrate not wet; 70-90 % is an estimate. Keep the land section damp without letting it turn to mud, and part of the substrate may well be drier.
WATER EN WATERKWALITEIT	The water section takes up roughly half of the enclosure. The depth is not documented for the species; 10-15 cm is borrowed from *Tylototriton verrucosus*.
VERLICHTING EN UVB	UVB is not necessary, and the species is most active at dusk and at night.
VOEDING	Adults are fed chopped earthworms, crickets and other prey of suitable size. Waxworms and other fatty larvae are popular, but should not be a staple. With a varied diet, vitamin and mineral supplements are normally not necessary.
GEDRAG EN OMGANG	The species is slow and clumsy. 2-3 animals can be kept together when there are enough hiding places for each animal.
NATUURLIJK VERSPREIDINGSGEBIED	China.