herpy

SALAMANDERS EN WATERSALAMANDERS

Wenxian Knobby Newt

Tylototriton wenxianensis

Tylototriton wenxianensis is a medium-sized, almost entirely black crocodile newt with orange toe tips and tail edge. It lives hidden in forest and scrub in the mountains of central China and enters the water only to breed; the eggs are laid on land. Little is known about its care, so this guide is based on close relatives.

Familie: Salamandridae **Mild huidgif**

KERNGEGEVENS

HET DIER

Volwassen grootte	**13-16 cm** totale lengte incl. staart
Levensduur	**10-20 jaar**
Actief	**In de schemering en ’s nachts**
Levenswijze	**Bodembewonend en semi-aquatisch**
Leefgebied	**Bergbos en moerasgebied**
Huidgif	**Mild** Was je handen na contact en houd je vingers uit de buurt van je ogen en mond
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Larven in het water**
Legplaats	**Op de grond**
Eieren per seizoen	**30-80**
Uitkomen	**20-36 dagen** bij 22-24 °C
Larvenperiode	**80-130 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Andere larven en kleine waterongewervelden (watervlooien, muggenlarven)**
Broedzorg	**Geen**

VERZORGING

Min. terrarium	**60 × 40 × 40 cm** L × B × H voor 3 volwassen dieren
Temperatuur overdag	**18-24 °C**
Temperatuur ’s nachts	**15-20 °C**
Luchtvochtigheid	**70-90%**
Water	**Watergedeelte** diepte 3-5 cm
UVB	**Niet nodig**
Voeding	**Insecten en wormen/slakken**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
EU-bijlage	**Bijlage B**
IUCN	**Kwetsbaar (VU)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Tylototriton wenxianensis*.
TEMPERATUUR	In summer the enclosure is kept at 22-24 °C by day, in winter at around 18 °C, and at night it may fall to 15-20 °C. Brief spells of lower temperatures do no harm.
LUCHTVOCHTIGHEID EN SPROEIEN	In summer the enclosure is misted daily so that the moss and substrate stay moist and the humidity is 70-90 %.
WATER EN WATERKWALITEIT	The water is soft and slightly acidic, pH 6.4-6.8, and at most about 5 cm deep. Outside the breeding season the animals hardly use the water section; in winter it can be almost dried out, leaving only a little water for humidity.
VERLICHTING EN UVB	UVB is not necessary. The species lives hidden and does not like strong light; the lighting is first and foremost for the plants. Provide about 12 hours of light in summer and 10-11 hours in winter.
VOEDING	The diet is not documented for the species. The genus takes ordinary newt food: earthworms, crickets, woodlice and waxworms. Adults are fed about twice a week, and in winter they eat irregularly and much less.
GEDRAG EN OMGANG	The animals live very secretively and can be kept alone, in pairs or in small groups. Keep the species on its own, without other amphibians. Handle the animals only when necessary, with clean, wet hands or powder-free gloves.
NATUURLIJK VERSPREIDINGSGEBIED	China.