herpy

SALAMANDERS EN WATERSALAMANDERS

Ziegler's Crocodile Newt

Tylototriton ziegleri

Tylototriton ziegleri is a small, almost entirely black crocodile newt of 13-16 cm with three rows of large warts along the back and clearly red fingers, toes and tail underside. The species was described in 2013, lives hidden in mountain forests in northern Vietnam and southern China and is easy to keep in an enclosure with a small water section.

Familie: Salamandridae Vroegere naam: *Yaotriton ziegleri* **Mild huidgif**

KERNGEGEVENS

HET DIER

Volwassen grootte	**13-16 cm** totale lengte incl. staart
Gewicht	**6-16 g**
Levensduur	**15-25 jaar**
Actief	**In de schemering en 's nachts**
Levenswijze	**Bodembewonend**
Leefgebied	**Bergbos**
Huidgif	**Mild** Was je handen na contact en houd je vingers uit de buurt van je ogen en mond
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Larven in het water**
Legplaats	**Op de grond**
Eieren per seizoen	**30-80**
Uitkomen	**20-36 dagen** bij 20-24 °C
Larvenperiode	**50-110 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Andere larven en kleine waterongewervelden (watervlooien, muggenlarven)**
Broedzorg	**Geen**

VERZORGING

Min. terrarium	**60 × 40 × 40 cm** L × B × H voor 3 volwassen dieren
Temperatuur overdag	**22-24 °C**
Temperatuur 's nachts	**20-22 °C**
Luchtvochtigheid	**80-100%**
Water	**Watergedeelte** diepte 1-5 cm
UVB	**Optioneel**
Winterrust	**16-18 °C** 12-17 weken
Voeding	**Insecten en wormen/slakken**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
EU-bijlage	**Bijlage B**
IUCN	**Kwetsbaar (VU)**

Verzorging in het kort

HET TERRARIUM
Tylototriton ziegleri is kept in a rainforest enclosure with a small water section. Three adults need an enclosure of at least 60 × 40 × 40 cm.

TEMPERATUUR
From March to October the enclosure is kept at 22-24 °C by day and 20-22 °C at night.

LUCHTVOCHTIGHEID EN SPROEIEN
Humidity is 80-100 %. The enclosure is misted daily with a plant sprayer or a rain system, and the amount is adjusted to the size and ventilation of the enclosure so that the substrate does not become stagnantly wet.

WATER EN WATERKWALITEIT
The water section must have soft, slightly acidic water with a pH of 6.4-6.8; for example rainwater that has stood for a while. The water is at most 5 cm deep. Never use tap water without removing the chlorine.

VERLICHTING EN UVB
Light is not particularly important to the animals, which do not like strong light. It is set to suit the plants: LED strips or fluorescent tubes for 12 hours in summer and 10-11 hours in winter. Indirect room light is also enough.

VOEDING
Adults are given food that is usual for newts: wax worms, crickets, house crickets, waxworm larvae, earthworms and isopods of suitable size. Adults are fed about twice a week; in winter they eat irregularly and much less.

GEDRAG EN OMGANG
The animals are hidden and quiet and are seen mostly when they are fed or when the males are in the water.

NATUURLIJK VERSPREIDINGSGEBIED
Vietnam en China.