herpy

LANDSCHILDPADDEN EN WATERSCHILDPADDEN

Painted Turtle

Chrysemys picta

Chrysemys picta is a diurnal, semi-aquatic turtle from North America with red and yellow stripes on the neck, legs and shell margin. The shell reaches 13-25 cm, and the animal can live for many decades. An adult requires a large, filtered tank with several hundred litres of water, a warm basking spot and strong UVB.

Familie: Emydidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**13-25 cm** totale lengte incl. staart
Gewicht	**300-500 g**
Levensduur	**25-40 jaar**
Actief	**Overdag**
Levenswijze	**Semi-aquatisch**
Leefgebied	**Moerasgebied en gematigd bos en grasland**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**130 × 65 × 60 cm** L × B × H voor 1 volwassen dier
Temperatuur overdag	**20-30 °C**
Zonneplek	**30-40 °C**
Temperatuur ’s nachts	**15-24 °C**
Luchtvochtigheid	**50-80%**
Water	**Aquaterrarium** diepte 25-35 cm
Watertemperatuur	**21-27 °C**
UVB	**Onmisbaar** Fergusonzone 3
Voeding	**Insecten, kant-en-klaar voer, planten, gewervelde dieren en wormen/slakken**
Gehouden	**Solitair of in groepen**

KWEEK

Voortplanting	**Eierleggend**
Legselgrootte	**4-15 eieren**
Legsels per jaar	**1-5**
Incubatie	**72-80 dagen** bij 25-30 °C

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Bijlage B**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Keep the animal in an aquarium, a large tank or a closed indoor pond, not in an ordinary enclosure.
TEMPERATUUR	The surface of the basking spot should be about 40 °C, measured with an infrared thermometer a couple of hours after the lamps are switched on, and the air above the basking spot 30-35 °C.
LUCHTVOCHTIGHEID EN WATER	The turtle's humidity is the water itself, so the relative humidity of the air is not documented in detail for the species and is not decisive; 50-80 % in the room is ample. What is decisive is the cleanliness of the water.
WATER EN WATERKWALITEIT	The water is the animal's living space, so it must be deep, clean and warm. The water should be at least twice as deep as the shell is wide, and 25-35 cm suits adults. The temperature is 24-27 °C (see Temperature).
VERLICHTING EN UVB	UVB is a necessity. The species belongs to Ferguson zone 3-4. The UV index at the surface of the basking spot should be around 3.0-4.0, and there may well be areas with a lower UV index so that the animal can choose.
VOEDING	*C. picta* is omnivorous. In the wild it eats aquatic plants, algae, aquatic insects, crustaceans, snails, small fish and carrion. The young animals are mostly carnivorous, and with age the diet becomes more plant-based.
GEDRAG EN OMGANG	*C. picta* is diurnal, shy and quick to flee into the water. It is not tame and does not thrive on contact, so it should be handled only when necessary, for example for health checks or cleaning.
NATUURLIJK VERSPREIDINGSGEBIED	Canada, Verenigde Staten en Mexico.