herpy

SKINKEN

Ocellated (Bronze) Skink

Chalcides ocellatus

Chalcides ocellatus is a lively, diurnal skink from the Mediterranean region, North Africa and the Middle East with a smooth, cylindrical body, small legs and pale-edged eyespots on the back. It swims through loose sand, gives birth to live young and is inquisitive and easy to watch. It does well in a dry enclosure with a sandy substrate, a warm basking spot and humid nights.

Familie: Scincidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**15-30 cm** totale lengte incl. staart
Gewicht	**22-39 g**
Levensduur	**10-20 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend en gravend**
Leefgebied	**Woestijn en halfwoestijn**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**90 × 45 × 30 cm** L × B × H voor 1 volwassen dier
Temperatuur overdag	**24-29 °C**
Zonneplek	**33-37 °C**
Temperatuur ’s nachts	**18-21 °C**
Luchtvochtigheid	**15-35%**
Water	**Waterbak**
UVB	**Aanbevolen** Fergusonzone 2
Voeding	**Insecten en kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**Solitair of in groepen**

KWEEK

Voortplanting	**Levendbarend**
Worpgrootte	**2-12 jongen**
Legsels per jaar	**1-2**
Draagtijd	**90-120 dagen**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

A single animal needs an enclosure of at least 90 × 45 × 30 cm, and larger is better. Floor area matters more than height, because the species mostly moves on the ground, but it also climbs, so branches and rocks give variety.

TEMPERATUUR

Chalcides ocellatus loves the sun and needs a clear temperature gradient. The basking spot should have a surface temperature of 43-49 °C, measured with an infrared thermometer, and the air temperature above it should be 33-37 °C.

LUCHTVOCHTIGHEID EN WATER

Humidity should follow the rhythm of nature: dry by day and humid at night. By day the enclosure should dry out to 15-35 %, and at night it may rise to 90-100 %.

VERLICHTING EN UVB

The species is diurnal and needs strong light and UVB. It belongs to Ferguson zone 2, and the UV index should be 2-3 at the basking spot and lower in the rest of the enclosure.

VOEDING

The species is first and foremost an insectivore and in the wild takes almost anything it can overpower: locusts, crickets, ants, beetles, woodlice, spiders, centipedes and insect larvae.

GEDRAG EN OMGANG

Chalcides ocellatus is active and inquisitive, but not an animal for regular handling: it is slippery and fast and disappears easily if it becomes frightened.

NATUURLIJK VERSPREIDINGSGEBIED

Italië, Malta, Griekenland, Cyprus, Turkije, Libanon, Syrië, Jordanië, Israël, Palestijnse gebieden, Irak, Iran, Saoedi-Arabië, Jemen, Oman, Verenigde Arabische Emiraten, Turkmenistan, Pakistan, Marokko, Westelijke Sahara, Mauritanië, Algerije, Tunesië, Libië, Egypte, Niger, Mali, Soedan en Tsjaad.