herpy

SKINKEN

South-western Crevice Skink

Egernia napoleonis

A medium-sized, flat-bodied, diurnal skink from the rocks and forests of south-western Australia. *Egernia napoleonis* hides in crevices under bark and stones and gives birth to live young. Almost no documented knowledge exists on keeping it in captivity, so this guide draws on the species in the wild and on the close relatives *Egernia striolata*, *Egernia cunninghami* and *Egernia kingii*.

Familie: Scincidae Vroegere naam: *Tiliqua napoleonis; Contundo napoleonis; Egernia bos*

KERNGEGEVENS

HET DIER

Volwassen grootte	**13-14 cm** kop-romplengte
Levensduur	**10-20 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend en rotsbewonend**
Leefgebied	**Rotsachtig terrein en gematigd bos en grasland**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**100 × 50 × 60 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**25-30 °C**
Zonneplek	**35-40 °C**
Temperatuur 's nachts	**14-20 °C**
Luchtvochtigheid	**40-60%**
Water	**Waterbak**
UVB	**Onmisbaar** Fergusonzone 3
Voeding	**Insecten, fruit en planten**
Gehouden	**Solitair of in paren**

KWEEK

Voortplanting	**Levendbarend**
Worpgrootte	**1-6 jongen**
Legsels per jaar	**1**
Draagtijd	**90-120 dagen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage III**
EU-bijlage	**Bijlage C**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Almost no documented knowledge exists on the care of *Egernia napoleonis*.
TEMPERATUUR	By day the air should be 25-30 °C with a clear gradient from the cool end to the warm end. Under a halogen spot lies a flat stone or branch whose surface should reach 35-40 °C, measured with an infrared thermometer.
LUCHTVOCHTIGHEID EN WATER	Humidity should be 40-60 %. Mist the enclosure lightly a couple of times a week, and keep the substrate slightly damp under one hide in the cool end, so the animal can choose. A dish of fresh water should always be available.
VERLICHTING EN UVB	*Egernia napoleonis* is diurnal and needs UVB for as long as the light is on. A UVB tube in a reflector over the basking area, covering about half the length of the enclosure, gives Ferguson Zone 3.
VOEDING	The diet is not documented for *Egernia napoleonis*, so feeding follows the close relatives: an omnivorous diet with plenty of insects and a good deal of plant material.
GEDRAG EN OMGANG	*Egernia napoleonis* is a diurnal skink that uses crevices as hides and climbs on bark and rocks. Close relatives in the same group, such as *Egernia striolata*, live in family groups.
NATUURLIJK VERSPREIDINGSGEBIED	Australië.