herpy

SKINKEN

Southern White's Skink

Liopholis whitii

A medium-sized, diurnal rock skink from cool temperate southern Australia and Tasmania. *Liopholis whitii* lives in rock crevices and burrows under stones, lives in small family groups and gives birth to live young. It needs an enclosure with many narrow hiding places, a warm basking spot, a cool night and patience when putting animals together.

Familie: Scincidae Vroegere naam: *Egernia whitii; Scincus whitii*

KERNGEGEVENS

HET DIER

Volwassen grootte	**8-10 cm** kop-romplengte
Levensduur	**10-15 jaar**
Actief	**Overdag**
Levenswijze	**Rotsbewonend en gravend**
Leefgebied	**Rotsachtig terrein en gematigd bos en grasland**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**90 × 45 × 45 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**20-26 °C**
Zonneplek	**30-35 °C**
Temperatuur 's nachts	**12-18 °C**
Luchtvochtigheid	**40-60%**
Water	**Waterbak**
UVB	**Aanbevolen** Fergusonzone 2
Voeding	**Insecten, fruit en planten**
Gehouden	**In paren of in groepen**

KWEEK

Voortplanting	**Levendbarend**
Geslachtsrijp	**24-36 maanden**
Worpgrootte	**1-4 jongen**
Legsels per jaar	**1**
Draagtijd	**100-140 dagen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage III**
EU-bijlage	**Bijlage C**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little is known about keeping *Liopholis whitii*. A single animal can be kept in 60 × 30 × 40 cm.
TEMPERATUUR	By day the air should be 20-26 °C, and under a lamp at one end a flat stone should reach 30-35 °C measured with an infrared thermometer. At night the temperature should drop to 12-18 °C.
LUCHTVOCHTIGHEID EN WATER	Humidity should be 40-60 %. Keep the substrate dry, but make part of it slightly damp under a stone or a piece of bark so the animals can choose. Do not mist daily.
VERLICHTING EN UVB	The species is diurnal and basks on rocks. A UVB tube in a reflector above the basking spot should provide UVB corresponding to Ferguson zone 2.
VOEDING	*Liopholis whitii* is omnivorous. Offer insects such as mealworms, crickets and cockroaches, and supplement with finely chopped herbs and a little fruit or berries.
GEDRAG EN OMGANG	The species is territorial towards strangers and in the wild lives in small family groups whose members know each other.
NATUURLIJK VERSPREIDINGSGEBIED	Australië.