herpy

SKINKEN

Red-Eyed Bush Crocodile Skink

Tribolonotus gracilis

Tribolonotus gracilis is a shy, crepuscular skink from the rainforest of northern New Guinea with a red-orange ring around the eyes and rows of pointed, backward-curving scales down the back that give it an armoured, almost dragon-like appearance. It lives hidden in damp forest litter, can make sounds and lays only one egg at a time. It is best suited as a display animal, not for handling.

Familie: Scincidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**16-20 cm** totale lengte incl. staart
Levensduur	**10 jaar**
Actief	**In de schemering en 's nachts**
Levenswijze	**Bodembewonend en gravend**
Leefgebied	**Regenwoud en moerasgebied**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**60 × 45 × 45 cm** L × B × H voor 1 volwassen dier
Temperatuur overdag	**24-27 °C**
Zonneplek	**27-28 °C**
Temperatuur 's nachts	**22-24 °C**
Luchtvochtigheid	**70-90%**
Water	**Waterbak**
UVB	**Aanbevolen** Fergusonzone 1
Voeding	**Insecten en wormen/slakken**
Gehouden	**Solitair of in paren**

KWEEK

Voortplanting	**Eierleggend**
Geslachtsrijp	**36-48 maanden**
Legselgrootte	**1 eieren**
Legsels per jaar	**1-6**
Incubatie	**65-85 dagen** bij 23-27 °C

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Bijlage D**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

A single animal needs a terrarium of at least 60 × 45 × 45 cm, and bigger is better. A pair needs at least 90 × 45 × 45 cm.

TEMPERATUUR

Tribolonotus gracilis is tropical, but does not bask in the wild. Keep the terrarium at 24-26 °C at the cool end and 27-28 °C in a warm hide, and let the temperature fall to 22-24 °C at night.

LUCHTVOCHTIGHEID EN WATER

The species tolerates drying out very badly. The humidity must be 70-80 % by day and higher at night, preferably about 90 %, and the substrate must always be damp.

VERLICHTING EN UVB

The species is active at dusk and at night. The animal can manage and breed without UVB when the food is supplemented with calcium and vitamin D3, but a weak UVB light gives the best results and is recommended.

VOEDING

The species is insectivorous. The staple diet can be crickets, cockroaches (e.g. *Blaptica dubia* and *Blaberus discoidalis*), mealworms, superworms, hornworms, silkworms, black soldier fly larvae, earthworms and snails.

GEDRAG EN OMGANG

Tribolonotus gracilis is an animal to watch and not for handling. When seized, it squeals, plays dead, tries to flee or drops its tail, and this stresses it.

NATUURLIJK VERSPREIDINGSGEBIED

Papoea-Nieuw-Guinea en Indonesië.