herpy

SKINKEN

Spiny Skink

Tribolonotus novaeguineae

Tribolonotus novaeguineae is a small, shy skink from the lowland rainforest of New Guinea with rows of pointed, backward-curving scales down the back. It lives hidden in damp forest litter, is most active at dusk and lays only one egg at a time. Little is known about the care of this particular species, and it is best suited as a display animal.

Familie: Scincidae Vroegere naam: *Zonurus novaeguineae*

KERNGEGEVENS

HET DIER

Volwassen grootte	**16-20 cm** totale lengte incl. staart
Levensduur	**10 jaar**
Actief	**In de schemering en 's nachts**
Levenswijze	**Bodembewonend en gravend**
Leefgebied	**Regenwoud en moerasgebied**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**60 × 45 × 45 cm** L × B × H voor 1 volwassen dier
Temperatuur overdag	**24-27 °C**
Zonneplek	**27-28 °C**
Temperatuur 's nachts	**22-24 °C**
Luchtvochtigheid	**70-90%**
Water	**Waterbak**
UVB	**Aanbevolen** Fergusonzone 1
Voeding	**Insecten en wormen/slakken**
Gehouden	**Solitair of in paren**

KWEEK

Voortplanting	**Eierleggend**
Geslachtsrijp	**36-48 maanden**
Legselgrootte	**1 eieren**
Legsels per jaar	**1-6**
Incubatie	**65-85 dagen** bij 23-27 °C

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Bijlage D**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

Little is known about the care of *Tribolonotus novaeguineae*. The species lives in the same kind of rainforest floor as *Tribolonotus gracilis* from northern New Guinea, has the same size and largely the same requirements.

TEMPERATUUR

The daytime temperature should be 24-27 °C, and a warm hide at 27-28 °C keeps the animal going, even though it does not bask. At night 22-24 °C is suitable.

LUCHTVOCHTIGHEID EN WATER

The humidity should be 70-90 %, highest at night, and the substrate must always be damp.

VERLICHTING EN UVB

The species is active at dusk and needs no strong basking spot, but 12 hours of daylight a day maintains its daily rhythm.

VOEDING

Tribolonotus novaeguineae is insectivorous. The staple diet is crickets, cockroaches, earthworms, mealworms and silkworms, and waxworms and superworms are given in smaller amounts.

GEDRAG EN OMGANG

Tribolonotus novaeguineae is an animal to watch and not for handling. When seized, it may make sounds, play dead, try to flee or drop its tail, and this stresses it.

NATUURLIJK VERSPREIDINGSGEBIED

Indonesië en Papoea-Nieuw-Guinea.