herpy

SLANGEN

Checkered Garter Snake

Thamnophis marcianus

Thamnophis marcianus is a medium-sized, mainly diurnal garter snake with a black checkerboard pattern on a green to olive-brown back. It lives in the south-western United States, Mexico and Central America, almost always near water, and grows to 45-105 cm. It is hardy and can be kept on thawed mice, but it is a selective feeder, may eat other snakes of its own species and escapes through the smallest gaps.

Familie: Colubridae

KERNGEGEVENS

HET DIER

Volwassen grootte	**45-105 cm** totale lengte incl. staart
Gewicht	**30-120 g**
Levensduur	**7-10 jaar**
Actief	**Overdag en 's nachts**
Levenswijze	**Bodembewonend en semi-aquatisch**
Leefgebied	**Halfwoestijn, savanne en moerasgebied**
Moeilijkheidsgraad	**Beginner**

VERZORGING

Min. terrarium	**100 × 50 × 50 cm** L × B × H voor 1 volwassen dier
Temperatuur overdag	**23-27 °C**
Zonneplek	**29-32 °C**
Temperatuur 's nachts	**18-24 °C**
Luchtvochtigheid	**35-60%**
Water	**Waterbak**
UVB	**Aanbevolen** Fergusonzone 2
Winterrust	**12-15 °C** 8-16 weken
Voeding	**Gewervelde dieren en wormen/slakken**
Gehouden	**Solitair**

KWEEK

Voortplanting	**Levendbarend**
Geslachtsrijp	**18-24 maanden**
Worpgrootte	**6-35 jongen**
Legsels per jaar	**1-2**
Draagtijd	**80-105 dagen**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little species-specific knowledge exists on the care of *Thamnophis marcianus*.
TEMPERATUUR	There must be a clear temperature gradient. The basking spot should be 29-32 °C measured on the surface, the cool end 23-27 °C and the night 18-24 °C.
LUCHTVOCHTIGHEID EN WATER	The humidity should be 35-60 %, highest at the cool end and at night. The enclosure should not be wet, because constant damp causes blisters on the belly and fungal infections.
VERLICHTING EN UVB	*Thamnophis marcianus* is diurnal and thrives in strong daylight. UVB is not strictly necessary, but it is recommended and often makes the snake more active.
VOEDING	In the wild *Thamnophis marcianus* eats frogs, toads, tadpoles, small fish and earthworms.
GEDRAG EN OMGANG	*Thamnophis marcianus* is quick and alert and usually becomes calm if it is handled regularly from a young age. Let new animals settle in for about two weeks, and only begin handling them when they are feeding reliably.
NATUURLIJK VERSPREIDINGSGEBIED	Verenigde Staten, Mexico, Belize, Honduras, Nicaragua en Costa Rica.