herpy

VARANEN

Pygmy Mulga Monitor

Varanus gilleni

A small, slender and diurnal tree climber from the interior desert regions of Australia. *Varanus gilleni* grows to only 35-40 cm and is one of the least demanding monitors, but it needs a very hot basking spot, a tall enclosure and a fixed rhythm with a cool winter.

Familie: Varanidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**35-40 cm** totale lengte incl. staart
Gewicht	**60-80 g**
Levensduur	**8-20 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend en bodembewonend**
Leefgebied	**Woestijn en halfwoestijn**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**60 × 60 × 90 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**26-30 °C**
Zonneplek	**40-60 °C**
Temperatuur ’s nachts	**18-20 °C**
Luchtvochtigheid	**40-60%**
UVB	**Aanbevolen** Fergusonzone 3
Winterrust	**18-22 °C** 6-10 weken
Voeding	**Insecten en gewervelde dieren**
Gehouden	**Solitair, in paren of één mannetje met meerdere vrouwtjes**

KWEEK

Voortplanting	**Eierleggend**
Legselgrootte	**1-7 eieren**
Incubatie	**84-131 dagen** bij 26-32 °C

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
EU-bijlage	**Bijlage B**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

For an adult pair the enclosure must be at least 60 × 60 × 90 cm (length × width × height), and larger is better. Height is important, because the species climbs.

TEMPERATUUR

Varanus gilleni needs a very hot basking spot. The surface temperature of the branch under the heat source must be 40-60 °C, preferably 50-60 °C, measured with an IR thermometer.

LUCHTVOCHTIGHEID EN WATER

The species comes from dry regions and does not need high humidity. Keep humidity at 40-60 %, with a dry enclosure as the starting point and a damper corner or deep substrate where the animal can retreat.

VERLICHTING EN UVB

The species is diurnal and basks in high heat.

VOEDING

Varanus gilleni eats mostly small arthropods: crickets, grasshoppers, cockroaches, mealworms, spiders and other insects. In the wild it also takes small lizards, especially geckos, and sometimes birds' eggs.

GEDRAG EN OMGANG

Varanus gilleni is among the calmest monitors. It is far less fast and quick-tempered than, for example, *V. acanthurus* and *V. tristis*. It seldom bites, and even animals that are a little skittish at first soon settle down.

NATUURLIJK VERSPREIDINGSGEBIED

Australië.