herpy

POISON DART FROGS

Blue-bellied Poison Frog

Andinobates minutus

Andinobates minutus is one of the smallest poison dart frogs, only 13-16 mm from snout to vent. It lives in the leaf litter of humid lowland forest from central Panama to the Pacific coast of Colombia, and the male carries the tadpoles up into bromeliads. The species is not collected for the pet trade and is hardly kept, so little is known about its care.

Family: Dendrobatidae Former name: *Ranitomeya minuta; Dendrobates minutus; Minyobates minutus* Published 16 November 2026

KEY FACTS

THE ANIMAL

Adult size	**13-16 mm** snout-vent length
Lifespan	**5-10 years**
Active	**By day**
Lifestyle	**Terrestrial**
Habitat	**Rainforest**
Skin toxins in the wild	**Mild**
Skin toxins in captivity	**None** Captive-bred animals. Wild-caught animals can keep their toxins for a long time
Difficulty	**Experienced**

CARE

Min. enclosure	**45 × 45 × 45 cm** L × W × H for 2 adults
Temperature, day	**22-25 °C**
Temperature, night	**19-22 °C**
Humidity	**80-100%**
Water	**No water dish**
UVB	**Optional** Ferguson zone 1
Diet	**Small invertebrates (springtails, fruit flies)**
Kept	**In pairs**

BREEDING

Reproduction	**Egg-laying**
Development	**Tadpoles carried to water**
Egg-laying site	**On the ground**
Eggs per clutch	**1-8**
Larval period	**60-80 days** from hatching to metamorphosis
Larval diet	**Complete diet**
Parental care	**Carries the tadpoles**

PROTECTION AND PAPERWORK

CITES	**Appendix II**
IUCN	**Least Concern (LC)**

Care at a glance

THE ENCLOSURE

Little is known about the care of *Andinobates minutus*. The care is based on the habitat: warm, humid lowland forest with rain for most of the year and a thick layer of leaf litter.

TEMPERATURE

The daytime temperature should be 22-25 °C and the night 19-22 °C. The figures are borrowed from *Ranitomeya imitator*, which thrives best at 21-24 °C and tolerates 18-27 °C.

HUMIDITY AND MISTING

Humidity should be 80-100 %. Spray the enclosure 1-2 times a day with rainwater or reverse osmosis water so that the leaf litter is kept damp, or use a rain system on a timer.

WATER AND WATER QUALITY

The adult frogs do not need a water bowl; they take up water through the skin from spraying and the damp leaf litter.

LIGHTING AND UVB

The species is diurnal and lives in shade on the forest floor. LED lighting for about 12 hours a day is enough for the plants. UVB is not necessary; a weak UVB light in Ferguson zone 1 can be used if there is plenty of shade.

FEEDING

In the wild *Andinobates minutus* lives on mites, ants, springtails and insect larvae.

BEHAVIOUR AND HANDLING

The species is diurnal and stays in the leaf litter and by the hiding places. The males call from fixed spots and defend their territory, so a pair is the way to keep them.

NATIVE RANGE

Colombia and Panama.