herpy

AGAMY

Boulenger's Pricklenape

Acanthosaura crucigera

Acanthosaura crucigera is a brown to greyish-green forest agama with a dark, cross-shaped mark on the nape, a spine behind each eye and a high crest of spikes on the neck. It lives in mountain forests from Myanmar and Thailand to northern Malaysia and Cambodia, reaches around 25 cm and uses both trunks and the forest floor. It needs a tall, humid and cool enclosure where the air never gets warmer than 28 °C.

Rodzina: Agamidae

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**25-26 cm** długość całkowita z ogonem
Długość życia	**5-10 lat**
Aktywność	**Za dnia**
Tryb życia	**Nadrzewny i naziemny**
Środowisko	**Las deszczowy i las górski**
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**100 × 50 × 120 cm** D × S × W dla 1 dorosłego osobnika
Temperatura w dzień	**22-26 °C**
Miejsce wygrzewania	**28-30 °C**
Temperatura w nocy	**18-22 °C**
Wilgotność	**70-90%**
Woda	**Część wodna**
UVB	**Zalecane** Strefa Fergusona 2
Pokarm	**Owady i dżdżownice i ślimaki**
Utrzymanie	**Pojedynczo lub w parach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Wielkość zniesienia	**6-12 jaj**
Inkubacja	**110-130 dni** przy 25-27 °C

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	The care of *Acanthosaura crucigera* is described in only a few sources.
TEMPERATURA	*Acanthosaura crucigera* tolerates heat poorly. The air in the enclosure should be 22-26 °C by day and must not get warmer than 28 °C. If it gets warmer for any length of time, there is a risk of heatstroke, stress and dehydration.
WILGOTNOŚĆ I WODA	Humidity should be 70-90 % around the clock, with the highest values at night and in the morning. Mist the enclosure thoroughly with soft water a couple of times a day, so that droplets hang on plants and walls.
OŚWIETLENIE I UVB	The species is active by day, but lives in the shade of the forest and needs relatively weak UVB. Use a T5 HO tube of 6 % (Ferguson zone 2) in a reflective fitting over the mesh, covering around half the length of the enclosure.
ŻYWIENIE	The species takes almost anything it can overpower: crickets, cockroaches, locusts, silkworms, black soldier fly larvae, earthworms and small slugs.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	*Acanthosaura crucigera* relies on its camouflage and often sits perfectly still on a trunk or on the floor. If it is spotted, it darkens, rises up on its front legs, spreads its throat pouch and opens its mouth.
NATURALNY ZASIĘG	Kambodża, Malezja, Mjanma (Birma) i Tajlandia.