herpy

AGAMY

Bibron's Agama

Agama bibronii

Agama bibronii is a medium-sized, diurnal rock agama from Morocco and the neighbouring parts of Algeria and Western Sahara. In the breeding season the male has a coppery orange head and a bluish purple body. The species needs a spacious, dry enclosure, a very warm basking spot, strong UVB and cool nights. Little is known about its care.

Rodzina: Agamidae Dawna nazwa: *Agama impalearis*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**25-30 cm** długość całkowita z ogonem
Długość życia	**10-15 lat**
Aktywność	**Za dnia**
Tryb życia	**Skalny i naziemny**
Środowisko	**Półpustynia i tereny skaliste**
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**120 × 60 × 60 cm** D × S × W dla 1 dorosłego osobnika
Temperatura w dzień	**24-32 °C**
Miejsce wygrzewania	**38-43 °C**
Temperatura w nocy	**16-20 °C**
Wilgotność	**30-50%**
Woda	**Miska z wodą**
UVB	**Niezbędne** Strefa Fergusona 3
Pokarm	**Owady i rośliny**
Utrzymanie	**Pojedynczo lub jeden samiec z kilkoma samicami**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Dojrzałość płciowa	**12-24 miesięcy**
Wielkość zniesienia	**5-23 jaj**
Zniesienia w roku	**1-2**
Inkubacja	**41-67 dni** przy 28-32 °C

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	Little is known about the care of *Agama bibronii*. The species is rare in captivity, but has been bred over several generations.
TEMPERATURA	*A. bibronii* needs a clear gradient. The basking spot should have a surface temperature of 38-43 °C, measured with an infrared thermometer. The rest of the enclosure is 24-32 °C.
WILGOTNOŚĆ I WODA	Humidity should be 30-50 %. *Agama bibronii* comes from arid to semi-arid regions, so the enclosure should be dry, with good ventilation. Mist the walls lightly one or two mornings a week; the animal readily drinks droplets.
OŚWIETLENIE I UVB	*A. bibronii* is diurnal and basks all day, so it needs strong light and UVB equivalent to Ferguson zone 3. This is borrowed from *Agama agama*, but suits the open, sunlit habitats of the species.
ŻYWIENIE	*Agama bibronii* is primarily an insectivore. Offer a varied diet of well-fed insects: cockroaches, crickets, locusts and, in smaller amounts, mealworms and superworms. Fatty insects such as waxworms are for treats only.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	*A. bibronii* is fast, wary and territorial. Males defend an area with threatening postures: they raise the body, bob with the forelimbs, flare the throat and lash with the tail. Never keep two males together.
NATURALNY ZASIĘG	Maroko, Sahara Zachodnia i Algieria.