herpy

DRZEWOŁAZY

Niceforo's Poison Frog

Ameerega ingeri

Ameerega ingeri is a small, diurnal poison dart frog of 20-27 mm from the foothills of the Andes in southern Colombia. The back is uniformly dark, the flanks have dark blue spots, and there are bright orange spots in the armpits and on the calves. The species is hardly ever kept, so care is based on the climate of its range and on related species.

Rodzina: Dendrobatidae Dawna nazwa: *Dendrobates ingeri; Epipedobates ingeri; Phyllobates ingeri*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**20-27 mm** długość od pyska do kloaki
Długość życia	**10-15 lat**
Aktywność	**Za dnia**
Tryb życia	**Naziemny**
Środowisko	**Las deszczowy**
Toksyny skórne na wolności	**Słabe**
Toksyny skórne w niewoli	**Brak** Zwierzęta z hodowli. Osobniki odłowione na wolności mogą przez długi czas zachować toksyny
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**60 × 40 × 40 cm** D × S × W dla 2 dorosłych osobników
Temperatura w dzień	**24-26 °C**
Temperatura w nocy	**20-22 °C**
Wilgotność	**70-100%**
Woda	**Miska z wodą** głębokość 1-2 cm
UVB	**Opcjonalne** Strefa Fergusona 1
Pokarm	**Małe bezkręgowce (skoczogonki, muszki owocowe)**
Utrzymanie	**W parach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Rozwój	**Kijanki noszone do wody**
Miejsce składania jaj	**Na ziemi**
Liczba jaj w zniesieniu	**4-16**
Czas do wyklucia	**10-22 dni**
Okres larwalny	**70-90 dni** od wyklucia do metamorfozy
Pokarm larw	**Glony i rośliny i gotowa karma**
Opieka nad potomstwem	**Nosi kijanki**

OCHRONA I FORMALNOŚCI

CITES	**Załącznik II**
IUCN	**Brak danych (DD)**

Hodowla w skrócie

TERRARIUM	Almost nothing is known about the care of *Ameerega ingeri*, and the species is hardly ever kept.
TEMPERATURA	Keep the enclosure at 24-26 °C by day, and let the temperature fall to 20-22 °C at night. The temperature must not exceed 28 °C, because frogs tolerate heat poorly.
WILGOTNOŚĆ I ZRASZANIE	Humidity should be 70-80 % in the middle of the day and close to 100 % in the morning and evening. A rain system or a fogger with a timer is easiest; otherwise mist two to three times a day with reverse-osmosis water or rainwater.
WODA I JAKOŚĆ WODY	Provide a shallow water dish with 1-2 cm of water, so that the frogs can easily climb out of it, and with stones or moss at the edge. Use reverse-osmosis water, rainwater or dechlorinated tap water, and change it often.
OŚWIETLENIE I UVB	The lighting is primarily for the plants: 10-12 hours a day with LED lighting with a plant-suitable spectrum. It is not documented that *Ameerega ingeri* needs UVB.
ŻYWIENIE	In the wild *Ameerega ingeri* lives mostly on small ants and other small arthropods from the leaf litter.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	*Ameerega ingeri* is diurnal and most active in the morning. It lives on the ground in the leaf litter, but likes to sleep on leaves some way above the ground, so plants with broad leaves are an advantage.
NATURALNY ZASIĘG	Kolumbia.