herpy

JASZCZURKI WŁAŚCIWE

Broadley's Rough-scaled Plated Lizard

Broadleysaurus major

Broadleysaurus major is a stout, armoured lizard from the savannas and rocky country of Africa, with thick, keeled scales in transverse bands, a blunt snout and a tail that makes up less than half of the total length. It is diurnal and inquisitive, but dislikes handling. It needs a large enclosure with deep burrowing substrate, warm basking spots, UVB and a diet of insects, plants and a little fruit.

Rodzina: Gerrhosauridae Dawna nazwa: *Gerrhosaurus major*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**30-60 cm** długość całkowita z ogonem
Długość życia	**10-15 lat**
Aktywność	**Za dnia**
Tryb życia	**Skalny, naziemny i kopiący nory**
Środowisko	**Sawanna i tereny skaliste**
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**120 × 60 × 60 cm** D × S × W dla 1 dorosłego osobnika
Temperatura w dzień	**23-26 °C**
Miejsce wygrzewania	**40-46 °C**
Temperatura w nocy	**15-22 °C**
Wilgotność	**40-60%**
Woda	**Miska z wodą**
UVB	**Niezbędne** Strefa Fergusona 3
Pokarm	**Owady, owoce i rośliny**
Utrzymanie	**Pojedynczo**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Wielkość zniesienia	**2-3 jaj**
Inkubacja	**110-145 dni** przy 28-30 °C

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM

An adult needs an enclosure of at least 120 × 60 × 60 cm (length × width × height). Bigger is better, because the species is very active and covers large areas in the wild.

TEMPERATURA

The basking spot should have a surface temperature of 40-46 °C, measured with an infrared thermometer. The cool end is 23-26 °C, and at night the temperature falls to 15-22 °C.

WILGOTNOŚĆ I WODA

Relative humidity in the enclosure is 40-60%, but the enclosure should have a dry area of 20-40% and a humid area of 50-80%. The air should be relatively dry, while the animal must have free access to a humid hide.

OŚWIETLENIE I UVB

The species is diurnal and requires UVB. Without UVB the animal becomes ill. The UVB tube should be a T5 HO of 10-14%, in a fitting that covers about half the length of the enclosure, on the same side as the heat bulbs.

ŻYWIENIE

The species is omnivorous and lives mostly on insects, but also eats plants, flowers and fruit, and adults live on a larger proportion of plant matter. Juveniles need more energy-rich insects and small animals.

ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM

B. major is active and inquisitive, but also nervous and easily frightened. Juveniles are particularly shy and defensive. The animal rarely bites, but lashes with its tail when grabbed.

NATURALNY ZASIĘG

Kenia, Tanzania, Mozambik, Malawi, Zambia, Etiopia, Erytrea, Dżibuti, Togo, Demokratyczna Republika Konga, Republika Południowej Afryki, Eswatini, Botswana, Zimbabwe, Sudan Południowy, Republika Środkowoafrykańska, Somalia, Uganda, Kamerun, Nigeria, Benin i Ghana.