herpy

JASZCZURKI WŁAŚCIWE

Erhard's Wall Lizard

Podarcis erhardii

Podarcis erhardii is a medium-sized, diurnal wall lizard from Greece and the surrounding countries, especially the Aegean islands. It reaches 15-23 cm including the tail, is grey, brown or greenish with very variable markings, and feeds on arthropods. The species is common in the wild, but little is known about its care.

Rodzina: Lacertidae

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**15-23 cm** długość całkowita z ogonem
Długość życia	**5-10 lat**
Aktywność	**Za dnia**
Tryb życia	**Skalny i naziemny**
Środowisko	**Tereny skaliste i lasy i łąki strefy umiarkowanej**
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**100 × 50 × 50 cm** D × S × W dla 2 dorosłych osobników
Temperatura w dzień	**22-30 °C**
Miejsce wygrzewania	**40-45 °C**
Temperatura w nocy	**14-18 °C**
Wilgotność	**40-60%**
Woda	**Miska z wodą**
UVB	**Zalecane** Strefa Fergusona 3
Zimowanie	**8-12 °C** 8-12 tygodni
Pokarm	**Owady i małe bezkręgowce (skoczogonki, muszki owocowe)**
Utrzymanie	**W parach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Wielkość zniesienia	**1-4 jaj**
Zniesienia w roku	**1-2**
Inkubacja	**35-45 dni** przy 26-28 °C

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	Little is known about the care of *Podarcis erhardii*.
TEMPERATURA	By day the temperature should be 22-30 °C, and the basking spot should be 40-45 °C at the surface, measured with an infrared thermometer.
WILGOTNOŚĆ I WODA	Humidity should be moderate: 40-60 % by day. The enclosure must be able to dry out completely, because still, humid air causes skin problems and fungus.
OŚWIETLENIE I UVB	*Podarcis erhardii* is diurnal and lives in open, sunlit terrain, so the enclosure needs strong heat and UVB light. The species probably belongs to Ferguson zone 3, with a UV index of about 3-5 at the basking spot.
ŻYWIENIE	*Podarcis erhardii* eats mostly small arthropods. The feeder insects can be crickets, small grasshoppers, flies, fruit flies, springtails, spiders and woodlice, and mealworms and wax worms only in small amounts.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	*Podarcis erhardii* is fast and alert and flees when disturbed. Males are territorial, and two males together will fight.
NATURALNY ZASIĘG	Grecja, Albania, Macedonia Północna i Bułgaria.