herpy

JASZCZURKI WŁAŚCIWE

Balkan Wall Lizard

Podarcis tauricus

Podarcis tauricus is a robust, diurnal, ground-dwelling lizard from south-eastern Europe and Asia Minor. It reaches 20-24 cm including the tail, is brown and green with two pale stripes along the side, and lives in open places with sand, grass and scattered vegetation. The species is common in the wild, but little is known about its care.

Rodzina: Lacertidae

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**20-24 cm** długość całkowita z ogonem
Długość życia	**5-10 lat**
Aktywność	**Za dnia**
Tryb życia	**Naziemny**
Środowisko	**Lasy i łąki strefy umiarkowanej**
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**100 × 50 × 50 cm** D × S × W dla 2 dorosłych osobników
Temperatura w dzień	**22-30 °C**
Miejsce wygrzewania	**40-45 °C**
Temperatura w nocy	**14-18 °C**
Wilgotność	**40-60%**
Woda	**Miska z wodą**
UVB	**Zalecane** Strefa Fergusona 3
Zimowanie	**8-12 °C** 8-12 tygodni
Pokarm	**Owady i małe bezkręgowce (skoczogonki, muszki owocowe)**
Utrzymanie	**W parach**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Wielkość zniesienia	**2-10 jaj**
Zniesienia w roku	**2**
Inkubacja	**35-45 dni** przy 26-28 °C

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM

Little is known about the care of *Podarcis tauricus*, so the figures for care, incubation time and lifespan are borrowed from *Podarcis muralis*.

TEMPERATURA

By day it should be 22-30 °C, and the basking spot should be 40-45 °C at the surface, measured with an infrared thermometer.

WILGOTNOŚĆ I WODA

Humidity should be moderate: 40-60% by day. The enclosure must be able to dry out completely, because stagnant, humid air causes skin problems and fungus.

OŚWIETLENIE I UVB

Podarcis tauricus is diurnal and lives in open, sunlit terrain, so the enclosure needs strong heat and UVB light. The species belongs to Ferguson zone 3, with a UV index of around 3-5 at the basking spot.

ŻYWIENIE

Podarcis tauricus eats small arthropods. Feeder insects can be crickets, small locusts, flies, fruit flies, springtails, spiders and woodlice, and mealworms and waxworms only in small amounts.

ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM

Podarcis tauricus is fast and alert, often basks in the open and flees when disturbed. Males form territories and defend them against other males, and two males together will fight.

NATURALNY ZASIĘG

Ukraina, Mołdawia, Rumunia, Węgry, Macedonia Północna, Bułgaria, Grecja, Albania i Turcja.