herpy

KAMELEONY

Boettger's Chameleon

Calumma boettgeri

Calumma boettgeri is one of the smallest chameleons, only 8-10 cm long, with a soft, round-oval rostral appendage and subdued colours in yellow, greenish brown and beige. It lives in rainforest in north-western Madagascar, including the islands of Nosy Be and Nosy Komba. The species needs warm, humid air with cooler nights and is for the experienced keeper. Its boundaries have been unclear.

Rodzina: Chamaeleonidae Dawna nazwa: *Chamaeleo boettgeri; Chamaeleon boettgeri; Chamaeleon macrorhinus*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**8-10 cm** długość całkowita z ogonem
Długość życia	**3-5 lat**
Aktywność	**Za dnia**
Tryb życia	**Nadrzewny**
Środowisko	**Las deszczowy**
Trudność	**Doświadczony**

HODOWLA

Min. terrarium	**40 × 40 × 60 cm** D × S × W dla 1 dorosłego osobnika
Temperatura w dzień	**22-27 °C**
Miejsce wygrzewania	**27-29 °C**
Temperatura w nocy	**16-20 °C**
Wilgotność	**70-100%**
UVB	**Zalecane** Strefa Fergusona 2
Pokarm	**Owady i małe bezkręgowce (skoczogonki, muszki owocowe)**
Utrzymanie	**Pojedynczo**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Dojrzałość płciowa	**9 miesięcy**
Wielkość zniesienia	**4-5 jaj**
Inkubacja	**90 dni** przy 22-24 °C

OCHRONA I FORMALNOŚCI

CITES	**Załącznik II**
Załącznik unijny	**Załącznik B**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	Little is known about the care of *Calumma boettgeri*. The climate below is based on measurements from the species' habitats.
TEMPERATURA	By day the temperature should be 22-27 °C, with a basking spot of 27-29 °C. The temperature must not exceed 29 °C on the warmest branches.
WILGOTNOŚĆ I WODA	Humidity must be high: 70-80 % for part of the day and higher at night, up to 100 % after misting.
OŚWIETLENIE I UVB	The species is diurnal and needs bright daylight for 12 hours a day. UVB is recommended. Aim for Ferguson zone 2, that is a UV index of about 2-3 on the highest branches, with a T5 HO UVB tube and plenty of shade.
ŻYWIENIE	*C. boettgeri* eats small insects: fruit flies, springtails, small crickets and small grasshoppers. The feeder insects must be small, because the animal itself is so small. Young are fed daily, adults every other day.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	The species is kept singly, because individuals are aggressive towards one another. Handling should be limited to what is necessary, and a new animal is given two weeks in peace.
NATURALNY ZASIĘG	Madagaskar.