herpy

KAMELEONY

Elliot's Chameleon

Trioceros ellioti

Trioceros ellioti is a small, live-bearing chameleon from montane forest, scrub and savanna in central and eastern Africa. Colours vary widely between populations: males often show blue and yellow tones, females are brown, grey or green, and both have a pale stripe along the flank. The species needs a cool enclosure with a clear night-time drop, high night humidity and solitary housing.

Rodzina: Chamaeleonidae Dawna nazwa: *Chamaeleo ellioti; Chamaeleo bequaerti*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**15-22 cm** długość całkowita z ogonem
Długość życia	**2-4 lat**
Aktywność	**Za dnia**
Tryb życia	**Nadrzewny**
Środowisko	**Sawanna i las górski**
Trudność	**Doświadczony**

HODOWLA

Min. terrarium	**45 × 45 × 90 cm** D × S × W dla 1 dorosłego osobnika
Temperatura w dzień	**20-24 °C**
Miejsce wygrzewania	**27-30 °C**
Temperatura w nocy	**10-15 °C**
Wilgotność	**30-100%**
UVB	**Niezbędne** Strefa Fergusona 3
Pokarm	**Owady**
Utrzymanie	**Pojedynczo**

ROZMNAŻANIE

Rozmnażanie	**Żyworodny**
Dojrzałość płciowa	**6-9 miesięcy**
Wielkość miotu	**2-18 młodych**
Zniesienia w roku	**1-4**
Ciąża	**90-160 dni**

OCHRONA I FORMALNOŚCI

CITES	**Załącznik II**
Załącznik unijny	**Załącznik B**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	A single adult needs an enclosure of at least 45 × 45 × 90 cm (length × width × height).
TEMPERATURA	By day the temperature should be 20-24 °C, preferably with a basking spot of 27-30 °C. The daytime temperature in the rest of the enclosure must not stay above 29 °C for long.
WILGOTNOŚĆ I WODA	Night is the humid time: 75-100 % at night, while humidity falls to 30-50 % by day.
OŚWIETLENIE I UVB	The species is diurnal and needs strong light. Four T5 HO tubes of 6500 K across the width of the enclosure light a 120 cm tall enclosure well.
ŻYWIENIE	*T. ellioti* is insectivorous and takes the usual feeder insects: crickets, grasshoppers, flies, cockroaches and silkworms. Flying insects are a favourite.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	The species is kept singly. It is milder than many chameleons, but reacts suddenly and violently when two animals meet, and can seriously injure an animal in the same enclosure.
NATURALNY ZASIĘG	Uganda, Rwanda, Burundi, Demokratyczna Republika Konga, Kenia, Sudan Południowy i Tanzania.

Pełny poradnik z mapą i aktualizacjami: herpy.org/en/chameleons/trioceros-ellioti