herpy

LEGWANY

Common Puerto Rican Anole

Anolis cristatellus

Anolis cristatellus is a stocky, diurnal anole from Puerto Rico and the Virgin Islands. The male has a tall, fin-like crest on the tail that is always erect, and a greenish-yellow dewlap with an orange-red margin, which he extends while bobbing and doing push-ups. The species lives on trunks and low down in the vegetation, is territorial and very shy, and needs UVB, live food and an enclosure with both branches and floor space.

Rodzina: Dactyloidae Dawna nazwa: *Anolis cozumelae; Anolis lindeni; Ctenonotus cristatellus*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**15-20 cm** długość całkowita z ogonem
Długość życia	**4-8 lat**
Aktywność	**Za dnia**
Tryb życia	**Nadrzewny i naziemny**
Środowisko	**Suchy las i las deszczowy**
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**60 × 45 × 60 cm** D × S × W dla 1 dorosłego osobnika
Temperatura w dzień	**26-29 °C**
Miejsce wygrzewania	**32-35 °C**
Temperatura w nocy	**20-24 °C**
Wilgotność	**60-80%**
Woda	**Miska z wodą**
UVB	**Niezbędne** Strefa Fergusona 2
Pokarm	**Owady**
Utrzymanie	**Pojedynczo lub jeden samiec z kilkoma samicami**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Wielkość zniesienia	**1-2 jaj**
Inkubacja	**45-55 dni** przy 26-28 °C

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	Little is known about keeping *Anolis cristatellus*.
TEMPERATURA	The air should be 26-29 °C by day, and the basking spot should have a surface temperature of 32-35 °C, measured with an infrared thermometer. The cooler side is 24-26 °C. At night 20-24 °C is suitable.
WILGOTNOŚĆ I WODA	Humidity should be 60-80 %: lower in the middle of the day and higher at night.
OŚWIETLENIE I UVB	*Anolis cristatellus* is diurnal and needs UVB every day to produce vitamin D3 and absorb calcium. The UVB lamp should give Ferguson zone 2, i.e. a moderate UV index at the basking spot.
ŻYWIENIE	The species is insectivorous and takes live prey that moves: crickets, cockroaches (such as dubia and discoid), grasshoppers, silkworms, hornworms, small superworms and soldier fly larvae.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	The species is shy, quick and not an animal for handling. It can drop its tail, and it does not grow back in the same form.
NATURALNY ZASIĘG	Portoryko, Wyspy Dziewicze Stanów Zjednoczonych i Brytyjskie Wyspy Dziewicze.

Pełny poradnik z mapą i aktualizacjami: herpy.org/en/iguanas/anolis-cristatellus