herpy

LEGWANY

Brown Anole

Anolis sagrei

Anolis sagrei is a small, fast, ground-oriented anole from Cuba, the Bahamas and the east coast of Central America. The male is brown with a yellow to deep red dewlap under the throat and defends his territory with head-bobs and push-ups. The species is hardy and active by day but very shy, and it needs UVB, live food and an enclosure with both branches and floor to run on.

Rodzina: Dactyloidae Dawna nazwa: *Anolis sagrae; Norops sagrei; Anolis ordinatus; Anolis greyi*

NAJWAŻNIEJSZE FAKTY

ZWIERZĘ

Wielkość dorosłego osobnika	**13-22 cm** długość całkowita z ogonem
Długość życia	**4-8 lat**
Aktywność	**Za dnia**
Tryb życia	**Nadrzewny i naziemny**
Środowisko	**Suchy las i las deszczowy**
Trudność	**Średnio zaawansowany**

HODOWLA

Min. terrarium	**60 × 45 × 60 cm** D × S × W dla 1 dorosłego osobnika
Temperatura w dzień	**26-29 °C**
Miejsce wygrzewania	**32-35 °C**
Temperatura w nocy	**20-24 °C**
Wilgotność	**60-80%**
Woda	**Miska z wodą**
UVB	**Niezbędne** Strefa Fergusona 2
Pokarm	**Owady**
Utrzymanie	**Pojedynczo lub jeden samiec z kilkoma samicami**

ROZMNAŻANIE

Rozmnażanie	**Jajorodny**
Dojrzałość płciowa	**9-12 miesięcy**
Wielkość zniesienia	**1-2 jaj**
Zniesienia w roku	**15-18**
Inkubacja	**28-56 dni** przy 26-28 °C

OCHRONA I FORMALNOŚCI

CITES	**Niewymieniony**
Załącznik unijny	**Niewymieniony**
IUCN	**Najmniejszej troski (LC)**

Hodowla w skrócie

TERRARIUM	A single adult needs an enclosure of at least 60 × 45 × 60 cm (length × width × height). Larger is better, and an enclosure of 60 × 60 × 90 cm gives far more to use without a much larger footprint.
TEMPERATURA	The air should be kept at 26-29 °C by day, and the basking spot should have a surface temperature of 32-35 °C measured with an infrared thermometer. The cooler side is 24-26 °C. At night 20-24 °C is suitable.
WILGOTNOŚĆ I WODA	Humidity should be 60-80 %: lower in the middle of the day and higher at night.
OŚWIETLENIE I UVB	*Anolis sagrei* is active by day and needs UVB every day to form D3 and absorb calcium. The UVB lamp should provide Ferguson zone 2, i.e. a UV index of at most about 3 at the basking spot.
ŻYWIENIE	The species is insectivorous and takes live prey that moves: crickets, cockroaches (such as dubia and discoid), grasshoppers, silkworms, hornworms, small superworms and soldier fly larvae.
ZACHOWANIE I OBCHODZENIE SIĘ ZE ZWIERZĘCIEM	The species is shy, fast and not an animal for handling. Captive-bred animals settle better than wild-caught ones.
NATURALNY ZASIĘG	Kuba, Bahamy, Kajmany, Jamajka, Meksyk, Belize, Gwatemala i Honduras.